科学通报 2026 年 第 71 卷 第 1 期: 234 ~ 251
评 述
《中国科学》杂志社
SCIENCE CHINA PRESS

非小细胞肺癌Wnt/β-catenin信号通路相关ncRNAs: 多维机制解析与诊疗转化研究

钟杨[1], 赖恒周[1], 何佳玮[1], 黄春霞[1], 杨澳羚[1], 黄文博[1], 马琼[1,3], 付西[1,2,3], 郑川[1*], 由凤鸣[1,2,3*]

1. 成都中医药大学附属医院, 代谢与慢病中医药防治四川省重点实验室, 成都 610075
2. 成都中医药大学肿瘤学教研室, 成都 610075
3. 成都中医药大学肿瘤研究所, 成都 610075
* 联系人, E-mail: zhengchuan@cdutcm.edu.cn; yfmdoc@163.com

2025-05-20 收稿, 2025-08-01 修回, 2025-09-19 接受, 2025-09-22 网络版发表
国家资助博士后研究人员计划(BX20250214)、国家自然科学基金(82405359)、中国博士后科学基金面上资助项目(2023MD744129)和成都市卫生健康委员会成都中医药大学委校联合创新基金(WXLH202403110)资助

摘要 Wnt/β-catenin信号通路是维持肺组织稳态的核心通路, 其异常激活则赋予肺癌恶性表型. 该通路复杂的调控机制已成为精准医疗的研究焦点. 大量研究表明, 非编码RNAs(non-coding RNAs, ncRNAs)是该通路的关键调控因子, 参与调控Wnt/β-catenin通路靶基因的表达, 在非小细胞肺癌(non-small-cell lung cancer, NSCLC)发生发展中扮演了重要角色. 本文通过系统解析不同类型ncRNAs在Wnt/β-catenin通路配体分泌、受体激活、信号转导及转录调控中的功能分工与协同网络, 揭示ncRNAs/Wnt/β-catenin轴驱动NSCLC发生发展的多维调控机制, 并进一步探讨了ncRNAs作为NSCLC早期诊断、疗效预测、预后评估的新型生物标志物以及开发为新型治疗靶点的临床诊疗转化研究进展与挑战. 本研究为深入理解NSCLC分子机制认知提供新视角, 并为推动其机制导向的精准诊疗转化奠定理论框架.

关键词 Wnt/β-catenin, ncRNAs, 非小细胞肺癌, 作用机制, 生物标志物

肺癌(lung cancer, LC)是全球癌症相关死亡的主要原因[1]. 非小细胞肺癌(non-small-cell lung cancer, NSCLC)是LC最常见的亚型, 约占全部LC病例的85%[2]. 对NSCLC分子机制认知不足、早期高灵敏/特异检测技术瓶颈及广泛耐药性仍是攻克NSCLC的主要障碍[3]. Wnt/β-catenin信号通路的异常激活已被证实是驱动NSCLC发生发展和治疗耐药的关键通路枢纽[4]. 近年研究发现, 以微小RNA(microRNA, miRNA)、长链非编码RNA(long non-coding RNA, lncRNA)和环状RNA(circular RNA, circRNA)为代表的非编码RNAs (non-coding RNAs, ncRNAs)通过靶向调控Wnt/β-catenin信号通路的配体、受体、信号转导分子、通路负调控因子及转录复合物层面, 在转录、转录后及表观遗传修饰水平影响NSCLC的恶性演进[5,6]. 然而, 目前不同类型ncRNAs在Wnt/β-catenin信号通路中的多水平、多层面的调控网络尚未系统解析, ncRNAs从基础发现到临床转化的系统路径亟待建立[7,8]. 本文系统阐述NSCLC发生发展中miRNA/lncRNA/circRNA调控Wnt/β-catenin通路的多层面分子调控网络, 展望靶向ncRNAs/Wnt/β-catenin轴的新型小分子药物及天然药物在临床ncRNAs药物开发中的前景, 并讨论Wnt/β-catenin通路相关ncRNAs作为NSCLC生物标志物的潜力, 以期为NSCLC的精准诊疗提供重要的理论基础和新方向(图1).

引用格式: 钟杨, 赖恒周, 何佳玮, 等. 非小细胞肺癌Wnt/β-catenin信号通路相关ncRNAs: 多维机制解析与诊疗转化研究. 科学通报, 2026, 71: 234–251
Zhong Y, Lai H Z, He J W, et al. NcRNAs in the Wnt/β-catenin signaling pathway: from multidimensional mechanisms to diagnostic and therapeutic translation in non-small-cell lung cancer (in Chinese). Chin Sci Bull, 2026, 71: 234–251, doi: 10.1360/CSB-2025-0706

© 2025《中国科学》杂志社



图 1 (网络版彩色)非小细胞肺癌Wnt/β-catenin通路中ncRNAs的层级化调控: 从机制到诊疗转化. 图片素材来源于BioRender.com

Figure 1 (Color online) Hierarchical regulation of ncRNAs in the Wnt/β-catenin pathway in NSCLC: from mechanisms to translational medicine. Created with BioRender.com

1 NSCLC中的Wnt/β-catenin信号通路

1.1 Wnt/β-catenin信号通路

Wnt/β-catenin信号通路又称作经典Wnt信号通路, 在包括发育、组织稳态和细胞增殖等各种生理过程中起着至关重要的作用[9]. 然而, 持续的Wnt/β-catenin信号通路激活赋予癌细胞持续的自我更新生长特性, 并与治疗耐药性相关[10]. 当Wnt/β-catenin信号通路处于失活状态时, 细胞质中两种支架蛋白AXIN(axis inhibition, AXIN)、APC(adenomatous polyposis coli, APC)以及两种激酶GSK-3β(glycogen synthase kinase 3 beta, GSK-3β)、CK-1α(casein kinase 1 alpha, CK-1α)所组成的β-catenin破坏复合物(β-catenin destruction complex, DC)诱导β-catenin磷酸化. 磷酸化的β-catenin通过与E3泛素连接酶β-TrCP(beta-transducin repeats-containing proteins, β-Trcp)结合而被泛素化, 最终被蛋白酶体(proteasome)降解, 维持胞内β-catenin的低水平状态[11,12]. 当Wnt/β-catenin通路处于激活状态时, WNT配体通过自分泌/旁分泌机制激活卷曲蛋白(frizzled, FZD)受体及低密度脂蛋白受体相关蛋白5/6(LDL-receptor-related protein 5/6, LRP5/6)共受体. 随后, FZD-LRP复合物磷酸化蓬乱蛋白(dishevelled, DVL), 募集支架蛋白AXIN和

APC到质膜上, 导致DC解离. 细胞质中积累的β-catenin经过细胞质-核穿梭, 与核内的T细胞因子/淋巴增强因子(T cell factor/lymphoid enhancer factor, TCF/LEF)结合[13], 激活*c-Myc*、*Cyclin D1*等促癌基因表达[14](图2).

1.2 NSCLC中Wnt/β-catenin信号通路异常激活的临床-分子特征

Wnt/β-catenin通路在正常肺组织中维持呼吸道上皮稳态与修复, 该通路异常激活是驱动NSCLC恶性演进的关键事件, 其复杂的调控机制已成为NSCLC精准医疗的研究焦点[5,15,16]. 越来越多的证据表明, Wnt/β-catenin信号通路组分与NSCLC临床-分子特征密切关联. WNT配体、β-catenin、FZD受体及通路负调控因子的表达水平与NSCLC患者预后、肿瘤亚型、肿瘤分期以及肿瘤微环境(tumor microenvironment, TME)中的免疫细胞浸润状态密切相关[17~20]. 此外, 异常的Wnt/β-catenin通路激活在NSCLC中广泛存在. 据估计, 大约35%~70%中的NSCLC具有异常的Wnt信号通路激活, 并且在76%的NSCLC中检测到异常β-catenin蛋白表达[21]. 多种WNT配体(如WNT1、WNT2B以及WNT5A)、FZD受体(如FZD2、FZD8)表达水平相较于癌旁正常组织显著上调[22~24], 并参与NSCLC的干性、增殖、血管生成、转移侵袭、治疗耐药性和其他关键肿瘤发展过程[25~28]. 然而, NSCLC中Wnt/β-catenin通路的激活模式不同于其他经典Wnt信号通路驱动的肿瘤. 多项研究表明NSCLC中的通路基因突变很少见[29,30], *APC*和*CTNNB1*基因特定位点突变在NSCLC中的发生率仅为10.4%和1.04%[31]. 这些证据表明, NSCLC中Wnt/β-catenin通路的激活并非主要依赖于高频驱动基因突变, 而是一个涉及多因素的复杂过程. 既存在部分基因的低频突变, 但更广泛地表现为通路组分的异常表达, 提示存在其他关键的调控机制.

2 ncRNAs: Wnt/β-catenin通路的多样性调控因子

ncRNAs是一类不编码蛋白质的功能性RNA分子, 其中miRNA、lncRNA和circRNA已被证明通过Wnt/β-catenin通路促进肿瘤发生、上皮-间充质转化(epithelial-mesenchymal transition, EMT)、血管生成、肿瘤发展、侵袭、转移及治疗耐药[32~34]. 不同类型ncRNAs调



图 2 (网络版彩色)Wnt/β-catenin信号通路. 图片素材来源于BioRender.com
Figure 2 (Color online) Wnt/β-catenin signaling pathway. Created with BioRender.com

控Wnt/β-catenin通路的分子模式存在显著差异.

2.1 miRNA/Wnt/β-catenin调控模式

miRNA的生物合成途径与作用机制在进化中高度保守. 其成熟依赖双链RNA前体经Drosha和Dicer (RNase III型酶)逐步加工, 进而与AGO(argonaute, AGO)蛋白结合形成RNA沉默复合体(RNA-induced silencing complex, RISC), 最终通过抑制靶mRNA翻译或促使其降解实现转录及转录后调控[35,36]. 如miR-200a既靶向EMT转录因子Zeb1/2, 又可直接阻断β-catenin mRNA翻译, 抑制Wnt/β-catenin通路激活, 其抑癌功能在胃腺癌和肝癌中得到进一步证实[37]. 此外, miRNA主要通过其5′端的种子序列与靶mRNA的3′非翻译区(3′ untranslated region, 3′UTR)进行部分碱基配对[38]. 这种不完全配对使miRNA可靶向多个mRNA. 例如miR-128-3p可同时靶向负调控AXIN1、分泌型卷曲相关蛋白2(secreted frizzled-related protein 2、SFRP2)和Wnt抑制因子1(Wnt inhibitory factor 1, WIF1)等多种通路关键组分发挥致癌作用[39](图3). 综上, 致癌/抑癌miRNA通过精准靶向Wnt/β-catenin通路核心元件, 以转录抑制、翻译阻断或mRNA降解等方式, 在肿瘤发生发展中起关键作用.

2.2 lncRNA/Wnt/β-catenin调控模式

lncRNA通过信号、支架、诱饵及引导等多种机制, 在转录、转录后、翻译后以及表观遗传水平参与染色质重塑、转录调控、生物分子凝聚物形成及蛋白质或RNA招募等过程[40,41]. lncRNA/Wnt/β-catenin信号转导贯穿肿瘤发生发展的各个阶段, 其作用模式具有高度多样性. 在转录水平, lncFZD6结合染色质重塑复合物SWI/SNF核心组分BRG1, 将其募集至FZD6启动子区, 激活肝癌起始细胞的Wnt/β-catenin通路[42]; 在转录后水平, lncRNA广泛作为竞争性内源RNA(competing endogenous RNA, ceRNA)结合特定miRNA, 解除



图 3 (网络版彩色)MiRNA生成及作用机制. Primary miRNA由RNA polymeraes II转录产生. Drosha将primary miRNA加工成precursor miRNA. 随后, precursor miRNA通过核转运蛋白进入细胞质, 与Dicer结合后被加工为成熟miRNA, 并装载至RISC复合体中, 其中一条链被降解, 另一条与RISC整合, 最终与靶mRNA互补结合并调控基因表达. lncRNA和circRNA也可与成熟的miRNA结合, 作为竞争性内源RNA抑制其活性. 图片素材来源于BioRender.com. AGO: argonaute蛋白; RISC: RNA沉默复合体

Figure 3 (Color online) MiRNA biogenesis and mechanism of action. Primary-miRNA is transcribed by RNA polymeraes II. Drosha processes primary-miRNA into precursor-miRNA. Subsequently, pre-miRNA is transported to the cytoplasm via nuclear transport proteins. In the cytoplasm, precursor-miRNA binds to Dicer and is processed into mature miRNA, which is loaded into RISC. One strand of the miRNA duplex is degraded, while the other integrates into RISC to ultimately bind complementary target mRNAs and regulate gene expression. Notably, both lncRNA and circRNA can also bind to mature miRNAs and act as competing endogenous RNAs to inhibit miRNA activity. Created with BioRender.com. AGO: argonaute; RISC: RNA-induced silencing complex

靶基因的表达抑制, 发挥转录后调控作用. 在翻译后水平, lncTIC1结合β-catenin N端结构域, 抑制GSK-3β介导的β-catenin磷酸化, 从而激活肝癌Wnt/β-catenin信号通路[43]. 在表观遗传水平, LINC01133通过诱导Wnt信号通路拮抗剂Dickkopf相关蛋白1(Dickkopf-1, DKK1)启动子甲基化, 激活Wnt/β-catenin通路, 促进胰腺癌进展[44](图4).

2.3 circRNA/Wnt/β-catenin调控模式

circRNA具有闭合环状结构, 主要定位于细胞质或外泌体中, 不易被RNA外切酶降解, 表达稳定性高[45,46]. 其功能机制与lncRNA类似: 一方面可作为miRNA海绵, 通过结合位点阻断miRNA与靶mRNA 3′UTR的相互作用, 解除下游基因的表达抑制[47]; 另一方面, circRNA通过结合并隔离特定蛋白干扰其功能; 或作为蛋白支架, 通过募集RNA结合蛋白(RNA-binding protein, RBP)或转录因子等, 进而调控复合体组装与蛋白活性[48,49]. 值得注意的是, 部分circRNA含内部核糖体进入位点, 可翻译为功能性多肽调控Wnt/β-catenin信号通路[50,51]. 例如, circAXIN1编码的AXIN1-295aa多肽竞争性结合APC破坏DC复合物结构[52], 而circβ-catenin则翻译生成β-catenin-370aa, 随后其作为分子诱饵竞争性拮抗GSK3β与β-catenin的结合, 从而抑制β-catenin磷酸化[53], 二者均通过干扰通路关键组分激活Wnt/β-catenin信号, 分别驱动胃癌与肝癌的恶性表型(图5).

3 ncRNAs作用于Wnt/β-catenin信号通路调控NSCLC发生发展

3.1 配体与受体层面的调控

3.1.1 靶向WNT的ncRNAs

WNT配体作为通路关键起始信号, 通过与细胞膜受体特异性结合, 激活下游Wnt/β-catenin通路传导, 从而调控细胞增殖、分化和稳态维持[54]. 在NSCLC发生发展中, ncRNAs通过直接或间接调控WNT配体的表达与功能, 成为Wnt/β-catenin通路异常激活的关键驱动因素[7,55,56]. 多项研究揭示了不同ncRNAs分别靶向WNT1、WNT2B及WNT5A等核心配体调控NSCLC恶



图 4 (网络版彩色)LncRNA作用机制. LncRNA可以通过信号、支架、诱饵和引导等作用机制参与细胞过程的各个方面. 图片素材来源于BioRender.com

Figure 4 (Color online) LncRNA mechanism of action. LncRNA can participate in various aspects of cellular processes through mechanisms of action such as signaling, scaffolding, decoy, and guiding. Created with BioRender.com



图 5 (网络版彩色)CircRNA作用机制. 图片素材来源于BioRender.com
Figure 5 (Color online) CircRNA mechanism of action. Created with BioRender.com

性进展. 其中miR-148a、miR-383可直接结合WNT1 mRNA 3′UTR区, 抑制Wnt/β-catenin通路活化, 降低NSCLC细胞迁移侵袭能力并诱导凋亡[57,58]. 在WNT2B调控中, miR-577通过下调WNT2B配体转录活性及蛋白表达水平, 抑制NSCLC细胞增殖和EMT过程[59]. 而WNT5A则表现出双重调控作用, 如miR-1253通过负调控WNT5A表达, 抑制NSCLC细胞增殖及转移[60]. 与之相反, miR-374a、circVAPA则通过靶向WNT5A, 促进NSCLC细胞EMT与干性维持[61,62]. 目前在哺乳动物中已发现19个独特的WNT配体并表现出组织特异性, 然而目前研究主要集中在少数几个WNT配体上. 针对其他多数WNT配体的特异性调控ncRNAs及其功能仍有待深入探索.

3.1.2 靶向FZD的ncRNAs

FZD受体构成Wnt/β-catenin通路的分子枢纽, 主导配体识别与跨膜信号转导过程[63]. ncRNAs对FZD家族成员的异常调控已被广泛证实[64,65], 并在Wnt/β-catenin通路的配体-受体互作层面表现出促癌或抑癌特性[66]. 在抑癌调控层面, 多种ncRNAs通过靶向FZD抑制Wnt/β-catenin通路激活[67]. 如miR-203直接结合FZD2 mRNA并阻断其翻译, 抑制Wnt/β-catenin通路活性[68]; lncRNA AK126698可通过负调控FZD8表达阻断通路信号转导, 诱导肿瘤细胞凋亡[69]; miR-3127-5p则通过抑制FZD4/β-catenin轴阻断NSCLC的EMT进程[70]. 与之相反, 促癌ncRNAs如circ_0017109吸附miR-671-5p, 解除其对FZD4 mRNA的翻译抑制, 进而激活Wnt/β-catenin通路并加速NSCLC进展[71]. 临床前研究表明, 针对FZD家族的靶向干预策略已展现出明确的抗肿瘤效应[72,73]. 系统阐释ncRNAs/FZD调控网络, 可能为NSCLC靶向治疗策略的创新提供分子基础.

3.1.3 靶向DVL的ncRNAs

DVL是一种支架蛋白, 帮助维持经典与非经典Wnt信号传导[74]. 尽管DVL蛋白是致癌Wnt/β-catenin通路的重要驱动因素, 然而DVL生物学活性调控仍知之甚少[75]. 近年来, ncRNAs对DVL磷酸化的调控机制逐渐阐明. 研究表明, LINC00673-v4在肺腺癌(lung adenocarcinoma, LUAD)中异常高表达, 其作为分子支架促进DDX3与CK1ε形成激酶复合体, 进而诱导DVL磷酸化, 最终激活Wnt/β-catenin通路, 驱动LUAD早期发生及转移[76]. 这一发现揭示了lncRNA通过激酶复合物调控DVL修饰的新致癌机制. 然而, DVL的泛素化、乙酰化等修饰机制及其生物学意义仍待阐明[77]. 新证据指向ncRNAs可能调控上述的翻译后修饰过程参与肿瘤等疾病进展[78,79], 全面解析DVL修饰机制有望拓展NSCLC靶向策略.

3.2 胞内信号转导节点的调控

3.2.1 直接靶向β-catenin的ncRNAs

ncRNAs通过ceRNA机制和翻译后修饰等方式直接调控β-catenin的表达与功能. 在ceRNA网络调控层

面，ncRNAs作为分子海绵作用解除靶miRNA对β-catenin mRNA的抑制，驱动NSCLC恶性进展．研究表明，lncRNA SNHG11既竞争性结合miR-4436a，解除CTNNB1 mRNA的翻译抑制，又直接结合β-catenin促进其核转位，展现出双重作用机制[80]．类似地，LINC01006[81]、lncRNA DANCR[82]以及lncRNA PKMYT1AR[83]均通过ceRNA模式激活Wnt/β-catenin通路．在翻译后修饰层面，circRNA通过编码功能肽调控β-catenin稳定性．其中，circFBXW7-185翻译为短肽circFBXW7-185AA，并以m6A依赖方式诱导β-catenin泛素化降解，进而逆转奥希替尼耐药及NSCLC干细胞特性[84]．已有研究表明circRNA或lncRNA编码的功能肽通过介导特定分子的磷酸化和泛素化，在多癌种中调控Wnt/β-catenin信号通路[85]．部分circRNA编码的肽或蛋白可作为肿瘤生物标志物，用于NSCLC的早期筛查与预后评估[86,87]．

3.2.2 间接靶向β-catenin的ncRNAs

同时，β-catenin的表达与功能也受到多种ncRNAs的多层级间接调控．在转录层面，抑癌因子miR-185通过抑制SRY-盒转录因子9(SRY-box transcription factor 9, SOX9)表达，下调β-catenin mRNA水平[88]，而lncRNA PVT1则通过吸附miR-361-3p解除对SOX9的抑制，诱导β-catenin转录激活，从而促进NSCLC增殖与侵袭[89]．在表观遗传层面，部分ncRNAs调控甲基化修饰抑制β-catenin功能，例如miR-708-5p通过抑制DNA甲基转移酶DNMT3A，降低细胞间黏附分子(E-Cadherin, CDH1)启动子甲基化，促进CDH1/β-catenin复合体形成，诱导Wnt/β-catenin通路失活[90]；lncRNA ITGB1-DT则通过竞争性结合增强子蛋白锌指同源物2，从而上调ITGB1表达并形成ITGB1/β-catenin/MYC正反馈回路，持续激活Wnt/β-catenin通路[91]．在翻译后修饰层面，ncRNAs调控乙酰化、泛素化修饰过程参与Wnt/β-catenin通路激活．如circCDR1as通过抑制富含丝氨酸/精氨酸的剪接因子1泛素化降解，促进β-catenin表达[92]；circRAC-GAP1则通过募集多聚嘧啶区结合蛋白1，并诱导下游的诱导复制时间调节因子1脱乙酰化，激活Wnt/β-catenin通路并促进CSC特性[93]．以上机制共同揭示了ncRNAs多水平、多途径调控网络在NSCLC进展中的核心作用．

3.2.3 靶向DC复合物的ncRNAs

DC复合物介导的β-catenin磷酸化和泛素化降解过程失调是Wnt/β-catenin通路过度激活的核心[94]，而ncRNAs通过干预DC组装与激酶活性，驱动NSCLC恶性进展．其中促癌因子miR-3607和miR-4326分别靶向APC及APC2的3′UTR区域阻断DC复合物组装，导致β-catenin核累积[95,96]；miR-17-92簇则通过抑制p38α激酶活性，间接促进GSK3β磷酸化，介导Wnt/β-catenin通路激活，以维持CSC特性[97]；同时，促癌因子miR-19[98]和miR-1246[99]直接抑制GSK3β表达，促进NSCLC细胞的CSC活性及EMT进程；另外，miR-20b可通过抑制APC表达，并形成正反馈循环加速NSCLC进展[100]．相反，部分ncRNAs靶向DC复合物发挥抑癌作用，如circ-GSK3B竞争结合FK506结合蛋白51，阻断GSK3β磷酸化，从而增强DC介导的β-catenin降解[101]；LINC00222则直接激活GSK3β，抑制Wnt/β-catenin通路活性，最终抑制LUAD细胞的增殖、迁移和CSC特性[102]．

3.3 靶向TCF/LEF转录复合物的ncRNAs

Wnt/β-catenin通路的终末效应依赖于β-catenin与核内TCF/LEF转录复合物的结合并激活下游靶基因转录，而这一关键过程受ncRNAs调控[103]．研究发现，在LUAD中，细胞核内的lncRNA UPLA1通过与桥粒斑蛋白(desmoplakin, DSP)结合，解除DSP对TCF/LEF转录活性的抑制，最终激活Wnt/β-catenin信号传导，促进LUAD的进展[104]．与之相反的是，抑癌因子chromobox蛋白同源物7(chromobox protein homolog 7, cbx7)通过阻断TCF4与β-catenin的结合抑制转录复合物形成，而lncRNA-SNHG7与miRNA-181相互作用，上调cbx7表达水平，进而破坏细胞核内的TCF4和β-catenin复合物形成，阻断Wnt/β-catenin通路激活，抑制LUAD细胞增殖和迁移[105]．可见，ncRNAs通过直接或间接干预TCF/LEF-β-catenin转录复合物的形成与活性，在Wnt/β-catenin通路中发挥关键作用．

3.4 靶向通路负调控因子的ncRNAs

Wnt/β-catenin通路的稳态由分泌型拮抗蛋白如DKK、SFRP、WIF家族与胞内调控元件共同维持[106]．ncRNAs通过直接或间接干预负调控因子表达驱动NSCLC恶性转化[107]．在直接调控方面，miR-1275通过抑制DKK3、SFRP1增强LUAD干细胞特性及转移[108]．同样，miR-582-3p通过抑制AXIN2、DKK3、SFRP1维持CSC特性[109]，miR-128-3p则同时靶向Axin1、SFRP2、WIF1等通路负调控因子，诱导CSC干性、EMT编程以及化疗耐药相关转移[39]．在间接调控方面，

LINC00326竞争结合miR-657，解除其对DKK2抑制作用，从而激活Wnt/β-catenin通路[110]；LINC01089则通过吸附miR-27a上调SFRP1，阻断Wnt/β-catenin通路传导[111]；此外，circ_0006427经miR-6783-3p/DKK1轴抑制LUAD增殖侵袭[112](图6)．这些发现表明ncRNAs通过多层面、多靶点调控通路负调控因子的表达，参与Wnt/β-catenin通路的稳态维持，为理解NSCLC发生发展的分子机制提供了新视角(表1)．

4 ncRNAs/WNT轴的治疗前景

4.1 ncRNAs/Wnt/β-catenin轴: NSCLC治疗耐药和靶向药物开发中的潜在靶点

Wnt/β-catenin通路相关ncRNAs在NSCLC治疗耐药中发挥核心调控作用[6]．目前越来越多的研究聚焦于ncRNAs介导的化疗抵抗机制，最新研究表明，hsa_circ_0125356通过miR-582-5p/FGF9轴调节WNT经典和非经典信号通路，诱导吉西他滨治疗耐药[113]．两种小分子拮抗剂WAY 316606和XAV-939可抑制hsa_circ_0125356诱导的Wnt/β-catenin通路的激活，为确定治疗NSCLC吉西他滨耐药的治疗靶点提供了新的方向[113]．此外，miR-130b通过抑制抑癌因子磷酸酶和张力蛋白同系物表达，降低β-catenin磷酸化水平，驱动Wnt/β-catenin通路介导的化疗耐药[114]．此外，lncRNA SNHG1通过miR-140-5p/Wnt轴促进顺铂(cisplatin, DDP)耐药[115]，而lncRNA RPPH1则通过miR-326/WNT2B轴加速NSCLC进展并诱导DDP抵抗[116]．这些发现系统揭示了ncRNAs通过不同分子机制调控Wnt/β-catenin通路驱动化疗耐药的共性特征，为开发新型联合治疗策略提供了理论依据．

在靶向药物开发方面，靶向ncRNAs为克服表皮生长因子受体酪氨酸激酶抑制剂(epidermal growth factor receptor-tyrosine kinase inhibitor, EGFR-TKI)耐药提供了新路径．多项研究揭示了ncRNAs的动态调控机制．



图 6 (网络版彩色)NSCLC中ncRNAs作为Wnt/β-catenin通路调控的分子开关. 图片素材来源于BioRender.com

Figure 6 (Color online) NcRNAs act as molecular switches governing Wnt/β-catenin signaling regulation in NSCLC. Created with BioRender.com

表 1 NSCLC中的ncRNAs/Wnt/β-catenin轴

Table 1 The ncRNAs/Wnt/β-catenin functional axis in NSCLC

作用位点	ncRNAs	细胞系	生物学功能	ncRNAs表达	参考文献
靶向WNT配体	miRNA-148a	NSCLC细胞系(A549、H1299)	抑制NSCLC细胞迁移和侵袭	下调	[57]
	miR-383	NSCLC细胞系(H1299、A549、和SW900)	抑制NSCLC细胞的细胞活力并诱导细胞凋亡	下调	[58]
	miR-577	NSCLC细胞系(H650、A549、H522、H1299和H1155)	抑制NSCLC细胞增殖和EMT过程	下调	[59]
	miR-1253	非小细胞细胞系(A549、NCI-H1299、NCI-H157、ANIP-973和GLC-82)	抑制NSCLC细胞的增殖、迁移和侵袭	下调	[60]
	miR-374a	NSCLC细胞系(HCC827和Calu1/HCC827-Gef抗性细胞系)	促进NSCLC细胞EMT、干性和吉非替尼耐药性	上调	[61]
	circVAPA	LC细胞系(H1299、H1975、A549和HCC827)	促进LC细胞增殖、迁移、侵袭和干性	上调	[62]
靶向FZD	miR-203	NSCLC细胞系h1299	抑制NSCLC细胞增殖、转移	下调	[68]
	lncRNA AK126698	NSCLC细胞系(A549、NCI-H520和H1299)	抑制NSCLC细胞增殖和迁移,诱导细胞凋亡	下调	[69]
	miR-3127-5p	LC细胞系(A549、H1299和HEK-293T)	抑制NSCLC中EMT, 细胞迁移、侵袭和黏附	下调	[70]
	circ_0017109	NSCLC细胞系(CALU3、CALU6、A549、H1229)	促进NSCLC细胞增殖、迁移、侵袭和干性	上调	[71]
靶向DVL	LINC00673-v4	LUAD细胞系(HCC827、NCI-H1650、A549、NCI-H596、NCI-H1975、NCI-H1299、SK-LU-1、NCI-H358、NCI-H2009、HCC4006、NCI-H2030和PC9)	促进LAD细胞侵袭、迁移和转移	上调	[76]
直接靶向β-catenin	lncRNA SNHG11	LUAD细胞系(A549、H1299、H460和SPCA1)	促进LUAD进展	上调	[80]
	LINC01006	LUAD细胞系(PC9、H1650、H1975和A549)	促进LUAD细胞增殖、迁移和EMT	上调	[81]
	lncRNA DANCR	LUAD细胞系(A549、H1975、H1755、H1944、H2087和H358NSCLC)大细胞癌细胞系(H661和H1299)	促进CSC的自我更新	上调	[82]
	lncRNA PKMYT1AR	LC细胞系(H358、H1975、H1299、H1650、A549和SPC-A1)	维持CSC特性	上调	[83]
	circFBXW7	奥希替尼耐药细胞系(HCC827OR和H1975OR)	抑制CSC特性, 逆转奥希替尼耐药	下调	[84]
间接靶向β-catenin	miR-185	NSCLC细胞系(A549、H1299、H1650和SK-MES-1)	抑制NSCLC细胞增殖和侵袭	下调	[88]
	lncRNA PVT1	NSCLC细胞系(H1650、H1975、H460和A549)	促进NSCLC细胞增殖、迁移、侵袭	上调	[89]
	miR-708-5p	LC细胞系(A549、Calu-3和95D)	抑制CSC特性	下调	[90]
	lncRNA ITGB1-DT	LUAD细胞系A549(ATCC CCL-185)、NCI-H1975(ATCC CRL-5908)	促进LUAD进展	上调	[91]
	circCDR1as	NSCLC细胞系(H1299、A549和H460)	促进NSCLC细胞增殖、迁移和侵袭	上调	[92]
	circRACGAP1	NSCLC细胞系(A549、H1299H1975、H460和PC-9)	维持CSC特性和转移	上调	[93]
靶向DC复合物	miR-3607	LC细胞系(NCI-H1650、5D、A549、NCI-H460、NCI-H358、NCI-H1975和NCI-H1299)	促进LC细胞增殖	上调	[95]
	miR-4326	LC细胞系(95D、NCI-H1299、NCI-H460、NCI-H1650、NCI-H358、NCI-H1975和A549)	促进LC细胞增殖	上调	[96]

(续表1)

作用位点	ncRNAs	细胞系	生物学功能	ncRNAs表达	参考文献
靶向DC复合物	miR-17-92	Lgr6+人肺干细胞	维持CSC特性	上调	[97]
	miR-19	NSCLC细胞系(A549和H1299)	促进CSC活性	上调	[98]
	miR-1246	NSCLC细胞系A549	促进EMT	上调	[99]
	miR-20b	LUAD细胞系(PC-9、H1975和A549) 非小细胞LC细胞系H1299	促进NSCLC细胞增殖、迁移和侵袭	上调	[100]
	circ-GSK3B	LUAD细胞系(PC9、H2073、H-1975和A549)	抑制LUAD细胞的增殖、迁移和干性	下调	[101]
	LINC00222	LUAD细胞系(SPC-A-1和LTEP-A-2)	抑制LUAD细胞迁移和侵袭	下调	[102]
靶向TCF/LEF转录复合物	LNCRNA UPLA1	LUAD细胞系(A549和H1299)	促进LUAD的迁移、侵袭	上调	[104]
	lncRNA-SNHG7	LUAD细胞系	抑制LUAD细胞增殖,抑制细胞凋亡	下调	[105]
靶向通路负调控因子	miR-1275	LC细胞系(L78、H460、A549、GLC-82、SPC-A1、PC9、H1299、H1975和H2228)	诱导LUAD细胞的干细胞样性状,促进肿瘤发生、复发和转移	上调	[108]
	miR-128-3p	LC细胞系(A549、Calu-3、SK-MES-1、PAa、NCI-H292、NCI-H520(H520)、NCI-H596、NCI-H1299、NCI-H1650、NCI-H1975、HLAMP和95D)	维持EMT和CSC编程以及诱导NSCLC细胞耐药	上调	[39]
	LINC00326	NSCLC细胞系(A549、NCI-H1299、HCC827、NCI-1650和NCI-H358)	抑制NSCLC进展	下调	[110]
	LINC01089	NSCLC细胞系	抑制NSCLC细胞的EMT	下调	[111]
	circ_0006427	LUAD细胞系(H1299、A549、SPC-A1和Calu3)	抑制LUAD进展	下调	[112]

例如, circ-FBXW7通过拮抗Wnt/β-catenin通路活性, 抑制CSC自我更新能力, 提示其作为逆转EGFR-TKI耐药的潜在靶点[84]; lncRNA NEAT1敲除可增强安罗替尼的促凋亡效应, 表明其可能成为联合治疗的增敏剂[117]; 此外, 外泌体miR-214通过激活Wnt/β-catenin通路, 赋予吉非替尼耐药性, 而靶向miR-214的antagomi可有效逆转耐药表型[118]. 值得注意的是, miR-374a与miR-548b通过协同作用调控NSCLC耐药细胞恶性表型, 前者通过抑制*WNT5a*阻断EMT及干细胞特性, 后者通过靶向*CCNB1*基因诱导细胞周期停滞, 这为多靶点干预策略提供了理论依据[119].

NSCLC治疗的复杂性不仅源于肿瘤细胞自身特性, 更与其所处的动态微环境密切相关, TME与ncRNAs/Wnt/β-catenin轴的调控网络正在成为突破现有治疗瓶颈的新方向[120,121]. 缺氧作为TME的核心特征, 可通过HIF-1α依赖性方式上调miR-1275, 协同激活Wnt/β-catenin通路, 显著增强LUAD干细胞特性并促进转移复发, 提示miR-1275可能成为治疗新靶点[108]. 在肿瘤细胞与TME的动态交流中, 外泌体介导的细胞间通讯发挥着关键作用. 研究发现外泌体转运的miR-4739同时激活肿瘤细胞及血管内皮细胞的Wnt/β-catenin通路, 促进"驱动基因阴性"NSCLC的恶性进展, 为开发针对此类难治性NSCLC的创新疗法提供了分子基础[122]. 另外, Wnt/β-catenin通路在肿瘤免疫微环境调控中同样具有重要作用. 研究表明, miR-23a/27a/24-2簇激活NSCLC中的β-catenin/TCF4轴驱动免疫逃逸以及PD-1/PD-L1抑制剂的耐药, 阻断该反馈回路可能是治疗miR-23a/27a/24-2簇中高表达的NSCLC的有用策略[123]. 针对ncRNAs/Wnt/TME轴的联合干预策略具有重要的临床转化价值, 有望突破当前NSCLC治疗的局限性.

4.2 临床ncRNAs药物开发及递送策略

RNA药物的靶向设计与递送系统开发是当前RNA疗法研究的核心热点及技术难点. 基于RNA疗法的临床转化可能受到与RNA药物靶向特异性、递送效率和体内耐受性等问题的阻碍, 但不可否认的是, 该领域为传统"不可成药"分子靶点及复杂疾病治疗开辟了独特机遇窗口[124]. 从分子优化层面而言, 针对RNA结构的化学修饰技术已取得显著进展, 通过对RNA分子的帽部结构、多聚腺苷酸尾部、核苷酸碱基(如假尿嘧啶替代尿嘧啶)及空间构象进行系统性修饰, 可有效提升

RNA稳定性、降低免疫原性并增强翻译效率[125~127]. 与此同时，基因编辑技术的革新为精准调控RNA功能提供了新维度[128]. CRISPR-Cas13系统通过特异性识别circRNA分子(如致癌性circFAM120A)实现精准编辑，而无需干扰同源线性mRNA，这为NSCLC相关ncRNAs干预提供了高特异性工具[129,130]. 在递送系统开发方面，外泌体、脂质纳米颗粒(lipid nanoparticle, LNP)、pH响应型聚合物载体等多元化递送平台已逐步成熟[131~133]. 已成功开发的各种纳米载体，例如外泌体、微泡、聚合物纳米颗粒、1,2-二油基-3-三甲基氯化铵纳米载体和病毒样颗粒在临床试验中已用于lncRNA递送[134]. 此外，将LNP与物理治疗技术(如离子照射)相结合，已显示出改善lncRNA治疗效果的前景，目前使用多孔氧化铁纳米剂递送的lncRNA CRYBG3已用于NSCLC治疗[135]. 值得关注的是，这些技术突破正加速RNA疗法的临床转化. 靶向KRAS G12D突变的小干扰RNA药物NDT-05-1040通过LNP实现肿瘤特异性递送[136]；基于miR-16模拟物的TargomiRs微型细胞则通过EGFR靶向机制在NSCLC临床试验中展现出可控安全性及初步疗效[137]. 然而我们必须清醒认识到，RNA疗法仍面临递送脱靶引发的正常组织毒性、载体免疫原性导致的全身炎症反应，以及TME异质性阻碍药物蓄积等多重挑战，这要求未来研究需进一步整合组织特异性配体修饰、动态生物传感等技术以完善治疗体系[138,139].

4.3 天然药物驱动ncRNAs靶向疗法

中医药是天然药物的重要组成部分，其核心特征在于通过多成分、多靶点、多通路协同调控机体. 这种整体调节模式与ncRNAs在细胞内形成的复杂调控网络高度契合. 大量研究发现，中药成分可通过多靶点、多成分干预ncRNAs的表达与功能，调控Wnt/β-catenin通路活性[140~142]. 例如，黄芩素不仅通过促进lncRNA-NEF表达，从而抑制Wnt/β-catenin信号通路激活，显著降低肿瘤细胞侵袭与转移能力，同时还可通过上调miR-25靶向沉默Wnt/β-catenin通路相关基因，上述作用共同拓展了黄芩素作为泛癌种潜在治疗剂的分子基础[143]. 进一步研究发现，黄酮类成分如黄芩苷通过激活miR-217/DKK1轴诱导结直肠癌细胞凋亡[144]，而川陈皮素通过抑制miR-15-5p，上调Wnt/β-catenin通路负调控因子AXIN2、WIF1表达，从而抑制NSCLC增殖[145]. 此外，天然多酚类化合物没食子酸可靶向lncRNA MALAT1阻断Wnt/β-catenin信号传导，抑制肿瘤进展[146]；萜类化合物如齐墩果酸则通过调控miR-122/β-catenin轴降低癌细胞迁移侵袭能力[147]，而芍药苷则通过miR-3194-5p/β-catenin轴抑制CSC特性[148]；青蒿琥酯则通过下调lncRNA-RP11，抑制Wnt/β-catenin通路活性，进而逆转肝癌细胞的EMT进程[149]. 已有的诸多研究从“天然产物-表观网络-核心通路”的级联调控模式系统揭示了中医药成分的多维作用机制，为开发基于天然产物的靶向治疗策略提供了理论依据[150].

5 Wnt/β-catenin通路相关ncRNAs的NSCLC生物标志物潜力

Wnt/β-catenin通路相关ncRNAs在NSCLC全周期管理中展现出多维度临床应用潜力，为NSCLC动态监测提供新型生物标志物. 首先，Wnt/β-catenin通路相关ncRNAs在NSCLC早期诊断及分型中展现出高灵敏性与特异性. 例如，血清外泌体miRNA组合(如miR-20b-5p和miR-3187-5p)可显著提高早期NSCLC检测效能(AUC=0.838)[151]，而LINC00673-v4联合miR-150可有效区分LUAD与肺鳞状细胞癌(AUC=0.784, P=0.03)[76]. 在预后评估方面，多项研究表明ncRNAs表达水平与患者生存显著相关. 例如，miR-582-3p高表达与较短的无复发生存期(recurrence free survival, RFS)及总生存期(overall Survival, OS)独立相关(P<0.01)[109]；LINC00222低表达提示LUAD复发风险升高(P<0.05)[102]，而miR-1275高表达患者OS显著降低且复发率增加，凸显其预后预警价值[108]. 在疗效评估方面，ncRNAs同样展现出重要价值. miR-128-3p高表达与基于DDP的NSCLC治疗疗效呈负相关，提示其作为疗效分层标志物的潜力[39]. 此外，ncRNAs表达谱可有效预测NSCLC的复发转移. 其中，miR-1275与β-catenin构建的联合预测模型具有优异的LUAD转移预测功能[108]；LINC00326、circ_0017109等ncRNAs与NSCLC临床分期及TNM(Tumor Node Metastasis, TNM)分类显著相关，为NSCLC复发监测提供分子基础[110,139]. 这些发现表明,Wnt/β-catenin通路相关ncRNAs通过其在诊断、预后和治疗预测等方面的应用价值，为NSCLC的精准诊疗提供了新的工具链.

6 总结与展望

随着精准医学的快速发展，ncRNAs动态分析技术为NSCLC诊疗模式的革新提供了重要契机，其不仅可

推动个体化治疗策略的制定，更可能通过早期诊断与实时监测显著改善患者预后. 然而, 当前研究多聚焦单一ncRNA分子或孤立通路节点制约了对ncRNAs调控网络的系统性认知. 本文系统解析ncRNAs/Wnt/β-catenin轴在NSCLC恶性演进中的多维调控机制，通过阐明不同类型ncRNAs(miRNA/lncRNA/circRNA)在通路配体分泌、膜表面受体激活、胞内信号转导及通路负调控因子的功能分工与协同效应, 揭示ncRNAs作为致癌/抑癌因子驱动NSCLC发生发展的复杂动态网络. ncRNAs作为Wnt/β-catenin通路的动态调控枢纽，展现出突破当前NSCLC治疗瓶颈的潜力. 基于此，靶向ncRNAs/Wnt/β-catenin轴可为NSCLC诊疗临床转化提供方向; 在诊断层面, Wnt/β-catenin通路相关ncRNAs展现出作为多维度生物标志物的潜力，其表达谱特征有望构建NSCLC“诊断-预后-疗效预测”全周期临床决策体系, 从而实现对NSCLC患者的精准分层管理; 在治疗策略方面, ncRNAs相关药物的开发及递送为NSCLC临床治疗打开了更为广阔的前景. 此外, 天然化合物以其多靶点特性，不仅为调控该复杂网络提供了丰富的可能性，并且基于天然化合物开发的靶向特定ncRNAs的小分子药物有望实现对信号通路的精准干预，从而提升新药研发效率, 推动NSCLC个体化治疗的发展.

当前关于ncRNAs在NSCLC中的研究仍存在重要挑战. 首先，研究结果呈现显著的异质性与矛盾性. 其主要原因在于不同临床样本、实验设计及检测技术等关键因素存在差异，从而导致ncRNAs作为生物标志物的可重复性受限. 基于多中心队列的大规模验证研究及标准化检测平台的建立是确证其临床有效性的必要前提. 其次，现有研究多聚焦于单个ncRNAs/Wnt/β-catenin轴的线性调控作用, 未能揭示ncRNAs在NSCLC信号网络中的全局调控特性. 最新研究表明，特定ncRNAs可同时调控Wnt、Hedgehog及Notch等通路关键节点，通过介导跨通路信号级联串扰驱动肿瘤恶性进展[152,153]. 未来，采用多组学整合策略系统解析ncRNAs在NSCLC信号网络中的多维调控图谱, 并建立以“分子机制解析-靶向治疗开发-标志物临床验证”为核心的临床诊疗转化路径，推动ncRNAs研究成果向临床实践的高效转化, 最终实现NSCLC诊疗的技术进步.

参考文献

1 Bray F, Laversanne M, Sung H, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin, 2024, 74: 229–263

2 Hendriks L E L, Remon J, Faivre-Finn C, et al. Non-small-cell lung cancer. Nat Rev Dis Primers, 2024, 10: 71

3 Melzer A C, Triplette M. Screening high risk populations for lung cancer. Br Med J, 2022, 376: o666

4 Stewart D J. Wnt signaling pathway in non-small cell lung cancer. J Natl Cancer Institute, 2013, 106: djt356

5 Zhong Y, He J W, Huang C X, et al. The NcRNA/Wnt axis in lung cancer: oncogenic mechanisms, remarkable indicators and therapeutic targets. J Transl Med, 2025, 23: 326

6 Lin S, Zhen Y, Guan Y, et al. Roles of Wnt/β-catenin signaling pathway regulatory long non-coding RNAs in the pathogenesis of non-small cell lung cancer. Cancer Manag Res, 2020, 12: 4181–4191

7 Xue C, Li G, Zheng Q, et al. The functional roles of the circRNA/Wnt axis in cancer. Mol Cancer, 2022, 21: 108

8 Braicu C, Zimta A A, Harangus A, et al. The function of non-coding RNAs in lung cancer tumorigenesis. Cancers, 2019, 11: 605

9 Jia Y, Liu Y, Li Y, et al. Persistent Wnt signaling affects IVF embryo implantation and offspring metabolism. Sci Bull, 2025, 70: 2297–2311

10 Rosenbluh J, Wang X, Hahn W C. Genomic insights into WNT/β-catenin signaling. Trends Pharmacol Sci, 2014, 35: 103–109

11 Xue C, Chu Q, Shi Q, et al. Wnt signaling pathways in biology and disease: mechanisms and therapeutic advances. Sig Transduct Target Ther, 2025, 10: 106

12 Wu X, Que H, Li Q, et al. Wnt/β-catenin mediated signaling pathways in cancer: recent advances, and applications in cancer therapy. Mol Cancer, 2025, 24: 171

13 Avşar E N, Çetin İ, Topçul M. An overview of the effect of the Wnt signaling pathway in lung cancer. Cell Mol Biol, 2022, 68: 41–46

14 Zhang Z, Westover D, Tang Z, et al. Wnt/β-catenin signaling in the development and therapeutic resistance of non-small cell lung cancer. J Transl Med, 2024, 22: 565

15 Rapp J, Jaromi L, Kvell K, et al. WNT signaling – lung cancer is no exception. Respir Res, 2017, 18: 167

16 Katoh M. Multi–layered prevention and treatment of chronic inflammation, organ fibrosis and cancer associated with canonical WNT/β–catenin signaling activation (Review). Int J Mol Med, 2018, 42: 713–725

17 Martín-Medina A, Cerón-Pisa N, Martinez-Font E, et al. TLR/WNT: a novel relationship in immunomodulation of lung cancer. Int J Mol Sci,

2022, 23: 6539
18 Krishnamurthy N, Kurzrock R. Targeting the Wnt/beta-catenin pathway in cancer: update on effectors and inhibitors. Cancer Treat Rev, 2018, 62: 50–60
19 Xue W, Cai L, Li S, et al. WNT ligands in non-small cell lung cancer: from pathogenesis to clinical practice. Discov Oncol, 2023, 14: 136
20 Raisch J, Côté-Biron A, Rivard N. A role for the WNT co-receptor LRP6 in pathogenesis and therapy of epithelial cancers. Cancers, 2019, 11: 1162
21 Xu X, Sun P L, Li J Z, et al. Aberrant Wnt1/β-catenin expression is an independent poor prognostic marker of non-small cell lung cancer after surgery. J Thoracic Oncol, 2011, 6: 716–724
22 Sumitomo R, Huang C L, Ando H, et al. Wnt2b and Wnt5a expression is highly associated with M2 TAMs in non−small cell lung cancer. Oncol Rep, 2022, 48: 189
23 Wang H, Xu M, Ma J, et al. Frizzled-8 as a putative therapeutic target in human lung cancer. Biochem Biophys Res Commun, 2012, 417: 62–66
24 Zhao H, Wu G, Luo Y, et al. WNT5B promotes the malignant phenotype of non-small cell lung cancer via the FZD3–DVL3–RAC1–PCP–JNK pathway. Cell Signal, 2024, 122: 111330
25 Malyla V, Paudel K R, De Rubis G, et al. Cigarette smoking induces lung cancer tumorigenesis via upregulation of the WNT/β-catenin signaling pathway. Life Sci, 2023, 326: 121787
26 Gujral T S, Chan M, Peshkin L, et al. A noncanonical Frizzled2 pathway regulates epithelial-mesenchymal transition and metastasis. Cell, 2014, 159: 844–856
27 Li J, Xu S, Dong H, et al. Altered Wnt5a expression affects radiosensitivity of non−small cell lung cancer via the Wnt/β−catenin pathway. Exp Ther Med, 2021, 23: 5
28 Entezari M, Deldar Abad Paskeh M, Orouei S, et al. Wnt/β-catenin signaling in lung cancer: association with proliferation, metastasis, and therapy resistance. Curr Cancer Drug Targets, 2024, 24: 94–113
29 Montpréville V T D, Ayoubi R E, Mamodaly M, et al. P1.09-17 CTNNB1 (beta-catenin) mutations in non-small cell lung carcinoma: a clinicopathological study of 18 cases. J Thorac Oncol, 2018, 13: s557
30 Pacheco-Pinedo E C, Morrisey E E. Wnt and Kras signaling-dark siblings in lung cancer. Oncotarget, 2011, 2: 569–574
31 Ros Montana F J, Iranzo P, Pedrola A, et al. WNT pathway mutations (APC/CTNNB1) and immune checkpoint inhibitors (ICI) response in metastatic non-small cell lung cancer (NSCLC) patients. Ann Oncol, 2019, 30: v799
32 Barati T, Mirzaei Z, Abarghooi Kahaki F, et al. Regulatory roles of ncRNAs in gastric cancer progression: a comprehensive insight into key signaling pathways. Med Oncol, 2025, 42: 309
33 Qiu W, Zhang S, Yu W, et al. Non-coding RNAs in hepatocellular carcinoma metastasis: remarkable indicators and potential oncogenic mechanism. Comput Biol Med, 2024, 180: 108867
34 Yu S, Han R, Gan R. The Wnt/β-catenin signalling pathway in haematological neoplasms. Biomark Res, 2022, 10: 74
35 Chen L L, Kim V N. Small and long non-coding RNAs: past, present, and future. Cell, 2024, 187: 6451–6485
36 Anastasiadou E, Jacob L S, Slack F J. Non-coding RNA networks in cancer. Nat Rev Cancer, 2017, 18: 5–18
37 Cong N, Du P, Zhang A, et al. Downregulated microRNA-200a promotes EMT and tumor growth through the Wnt/β-catenin pathway by targeting the E-cadherin repressors ZEB1/ZEB2 in gastric adenocarcinoma. Oncol Rep, 2013, 29: 1579–1587
38 Saw P E, Xu X, Chen J, et al. Non-coding RNAs: the new central dogma of cancer biology. Sci China Life Sci, 2021, 64: 22–50
39 Cai J, Fang L, Huang Y, et al. Simultaneous overactivation of Wnt/β-catenin and TGFβ signalling by miR-128-3p confers chemoresistance-associated metastasis in NSCLC. Nat Commun, 2017, 8: 15870
40 Mattick J S, Amaral P P, Carninci P, et al. Long non-coding RNAs: definitions, functions, challenges and recommendations. Nat Rev Mol Cell Biol, 2023, 24: 430–447
41 Ransohoff J D, Wei Y, Khavari P A. The functions and unique features of long intergenic non-coding RNA. Nat Rev Mol Cell Biol, 2017, 19: 143–157
42 Chen Z, Gao Y, Yao L, et al. LncFZD6 initiates Wnt/β-catenin and liver TIC self-renewal through BRG1-mediated FZD6 transcriptional activation. Oncogene, 2018, 37: 3098–3112
43 Chen Z, Yao L, Liu Y, et al. LncTIC1 interacts with β-catenin to drive liver TIC self-renewal and liver tumorigenesis. Cancer Lett, 2018, 430: 88–96
44 Weng Y C, Ma J, Zhang J, et al. Long non-coding RNA LINC01133 silencing exerts antioncogenic effect in pancreatic cancer through the methylation of DKK1 promoter and the activation of Wnt signaling pathway. Cancer Biol Ther, 2019, 20: 368–380
45 Zhang F, Jiang J, Qian H, et al. Exosomal circRNA: emerging insights into cancer progression and clinical application potential. J Hematol Oncol, 2023, 16: 67
46 Zhang J, Luo Z, Zheng Y, et al. CircRNA as an Achilles heel of cancer: characterization, biomarker and therapeutic modalities. J Transl Med,

2024, 22: 752
47 Chen W, Xu J, Wu Y, et al. The potential role and mechanism of circRNA/miRNA axis in cholesterol synthesis. Int J Biol Sci, 2023, 19: 2879–2896
48 Chen L, Shan G. CircRNA in cancer: fundamental mechanism and clinical potential. Cancer Lett, 2021, 505: 49–57
49 Chen J, Gu J, Tang M, et al. Regulation of cancer progression by circRNA and functional proteins. J Cell Physiol, 2022, 237: 373–388
50 Margvelani G, Maquera K A A, Welden J R, et al. Translation of circular RNAs. Nucleic Acids Res, 2025, 53: gkae1167
51 Yi Q, Feng J, Lan W, et al. CircRNA and lncRNA-encoded peptide in diseases, an update review. Mol Cancer, 2024, 23: 214
52 Peng Y, Xu Y, Zhang X, et al. A novel protein AXIN1-295aa encoded by circAXIN1 activates the Wnt/β-catenin signaling pathway to promote gastric cancer progression. Mol Cancer, 2021, 20: 158
53 Liang W C, Wong C W, Liang P P, et al. Translation of the circular RNA circβ-catenin promotes liver cancer cell growth through activation of the Wnt pathway. Genome Biol, 2019, 20: 84
54 Ash C, Purnell B A. How Wnt ligands achieve specificity. Science, 2018, 361: 657.12–659
55 Meng S, Zhou H, Feng Z, et al. CircRNA: functions and properties of a novel potential biomarker for cancer. Mol Cancer, 2017, 16: 94
56 Peng Y, Zhang X, Feng X, et al. The crosstalk between microRNAs and the Wnt/β-catenin signaling pathway in cancer. Oncotarget, 2017, 8: 14089–14106
57 Chen Y, Min L, Ren C, et al. miRNA-148a serves as a prognostic factor and suppresses migration and invasion through Wnt1 in non-small cell lung cancer. PLoS One, 2017, 12: e0171751
58 Gu B, Wang J, Song Y, et al. microRNA-383 regulates cell viability and apoptosis by mediating Wnt/β-catenin signaling pathway in non–small cell lung cancer. J Cell Biochem, 2019, 120: 7918–7926
59 Wang B, Sun L, Li J, et al. miR-577 suppresses cell proliferation and epithelial-mesenchymal transition by regulating the WNT2B mediated Wnt/β-catenin pathway in non-small cell lung cancer. Mol Med Report, 2018, 18: 2753–2761
60 Liu M, Zhang Y, Zhang J, et al. MicroRNA-1253 suppresses cell proliferation and invasion of non-small-cell lung carcinoma by targeting WNT5A. Cell Death Dis, 2018, 9: 189
61 Võsa U, Vooder T, Kolde R, et al. Identification of miR-374a as a prognostic marker for survival in patients with early-stage nonsmall cell lung cancer. Genes Chromosomes Cancer, 2011, 50: 812–822
62 Zhao Y, Dai Q, Fu X, et al. CircVAPA exerts oncogenic property in non-small cell lung cancer by the miR-876-5p/WNT5A axis. J Gene Med, 2021, 23: e3325
63 Sun M, Zhu H, Yu Y, et al. Role of the Wnt signaling pathway in the complex microenvironment of breast cancer and prospects for therapeutic potential (Review). Int J Oncol, 2025, 66: 1–22
64 Liu H, Sun X, Xiu S, et al. Frizzled receptors (FZDs) in Wnt signaling: potential therapeutic targets for human cancers. Acta Pharmacol Sin, 2024, 45: 1556–1570
65 Ueno K, Hirata H, Hinoda Y, et al. Frizzled homolog proteins, microRNAs and Wnt signaling in cancer. Intl J Cancer, 2012, 132: 1731–1740
66 DeBruine Z J, Xu H E, Melcher K. Assembly and architecture of the Wnt/β-catenin signalosome at the membrane. Br J Pharmacol, 2017, 174: 4564–4574
67 Sharma A, Mir R, Galande S. Epigenetic regulation of the Wnt/β-catenin signaling pathway in cancer. Front Genet, 2021, 12: 681053
68 Mine M, Yamaguchi K, Sugiura T, et al. miR-203 inhibits Frizzled-2 expression via CD82/KAI1 expression in human lung carcinoma cells. PLoS One, 2015, 10: e0131350
69 Fu X, Li H, Liu C, et al. Long noncoding RNA AK126698 inhibits proliferation and migration of non-small cell lung cancer cells by targeting Frizzled-8 and suppressing Wnt/β-catenin signaling pathway. Onco Targets Ther, 2016, 9: 3815–3827
70 Yang Y, Sun Y, Wu Y, et al. Downregulation of miR-3127-5p promotes epithelial-mesenchymal transition via FZD4 regulation of Wnt/β-catenin signaling in non-small-cell lung cancer. Mol Carcinog, 2018, 57: 842–853
71 Yang B, Zhang B, Qi Q, et al. CircRNA has_circ_0017109 promotes lung tumor progression via activation of Wnt/β-catenin signaling due to modulating miR-671-5p/FZD4 axis. BMC Pulm Med, 2022, 22: 443
72 Do M, Wu C C N, Sonavane P R, et al. A FZD7-specific antibody–drug conjugate induces ovarian tumor regression in preclinical models. Mol Cancer Ther, 2021, 21: 113–124
73 Do M, Wu C C, Adams S, et al. Abstract 1848: targeting FZD7-positive cancers using a novel antibody-drug conjugate. Cancer Res, 2021, 81: 1848
74 Bowin C F, Kozielewicz P, Grätz L, et al. WNT stimulation induces dynamic conformational changes in the Frizzled-dishevelled interaction. Sci Signal, 2023, 16: eabo4974
75 Shi Q, Chen Y G. Regulation of dishevelled protein activity and stability by post-translational modifications and autophagy. Trends Biochem Sci, 2021, 46: 1003–1016

76 Guan H, Zhu T, Wu S, et al. Long noncoding RNA LINC00673-v4 promotes aggressiveness of lung adenocarcinoma via activating WNT/β-catenin signaling. Proc Natl Acad Sci USA, 2019, 116: 14019–14028

77 Qin K, Yu M, Fan J, et al. Canonical and noncanonical Wnt signaling: multilayered mediators, signaling mechanisms and major signaling crosstalk. Genes Dis, 2024, 11: 103–134

78 Ryu H Y, Zhao D, Li J, et al. Histone sumoylation promotes Set3 histone-deacetylase complex-mediated transcriptional regulation. Nucleic Acids Res, 2020, 48: 12151–12168

79 Qi X, Wang H, Xia L, et al. miR-30b-5p releases HMGB1 via UBE2D2/KAT2B/HMGB1 pathway to promote pro-inflammatory polarization and recruitment of macrophages. Atherosclerosis, 2021, 324: 38–45

80 Liu S, Yang N, Wang L, et al. lncRNA SNHG11 promotes lung cancer cell proliferation and migration via activation of Wnt/β-catenin signaling pathway. J Cell Physiol, 2020, 235: 7541–7553

81 Zhang Y, Liu H, Zhang Q, et al. Long noncoding RNA LINC01006 facilitates cell proliferation, migration, and epithelial-mesenchymal transition in lung adenocarcinoma via targeting the microRNA 129-2-3p/CTNNB1 axis and activating Wnt/β-catenin signaling pathway. Mol Cell Biol, 2021, 41: e00380-20

82 Yu J E, Ju J A, Musacchio N, et al. Long noncoding RNA DANCR activates Wnt/β-catenin signaling through MiR-216a inhibition in non-small cell lung cancer. Biomolecules, 2020, 10: 1646

83 He Y, Jiang X, Duan L, et al. LncRNA PKMYT1AR promotes cancer stem cell maintenance in non-small cell lung cancer via activating Wnt signaling pathway. Mol Cancer, 2021, 20: 156

84 Li K, Peng Z Y, Wang R, et al. Enhancement of TKI sensitivity in lung adenocarcinoma through m^6A-dependent translational repression of Wnt signaling by circ-FBXW7. Mol Cancer, 2023, 22: 103

85 Wu P, Mo Y, Peng M, et al. Emerging role of tumor-related functional peptides encoded by lncRNA and circRNA. Mol Cancer, 2020, 19: 22

86 Meng E, Deng J, Jiang R, et al. CircRNA-encoded peptides or proteins as new players in digestive system neoplasms. Front Oncol, 2022, 12: 944159

87 Setrerrahmane S, Li M, Zoghbi A, et al. Cancer-related micropeptides encoded by ncRNAs: promising drug targets and prognostic biomarkers. Cancer Lett, 2022, 547: 215723

88 Lei Z, Shi H, Li W, et al. miR−185 inhibits non−small cell lung cancer cell proliferation and invasion through targeting of SOX9 and regulation of Wnt signaling. Mol Med Report, 2017, 27: 1015–1023

89 Qi G, Li L. Long non-coding RNA PVT1 contributes to cell growth and metastasis in non-small-cell lung cancer by regulating miR-361-3p/SOX9 axis and activating Wnt/β-catenin signaling pathway. Biomed Pharmacother, 2020, 126: 110100

90 Liu T, Wu X, Chen T, et al. Downregulation of DNMT3A by miR-708-5p inhibits lung cancer stem cell–like phenotypes through repressing Wnt/β-catenin signaling. Clin Cancer Res, 2018, 24: 1748–1760

91 Chang R, Xiao X, Fu Y, et al. ITGB1-DT facilitates lung adenocarcinoma progression via forming a positive feedback loop with ITGB1/Wnt/β-catenin/MYC. Front Cell Dev Biol, 2021, 9: 631259

92 Xu J, Huang L, Bao T, et al. CircCDR1as mediates $PM_{2.5}$-induced lung cancer progression by binding to SRSF1. Ecotoxicol Environ Saf, 2023, 249: 114367

93 Xiong H, Liu B, Liu X Y, et al. circ_rac GTPase–activating protein 1 facilitates stemness and metastasis of non–small cell lung cancer via polypyrimidine tract–binding protein 1 recruitment to promote sirtuin-3–mediated replication timing regulatory factor 1 deacetylation. Lab Invest, 2023, 103: 100010

94 Jung Y S, Park J I. Wnt signaling in cancer: therapeutic targeting of Wnt signaling beyond β-catenin and the destruction complex. Exp Mol Med, 2020, 52: 183–191

95 Lin Y, Gu Q, Sun Z, et al. Upregulation of miR-3607 promotes lung adenocarcinoma proliferation by suppressing APC expression. Biomed Pharmacother, 2017, 95: 497–503

96 Xu G, Zhang Z, Zhang L, et al. miR-4326 promotes lung cancer cell proliferation through targeting tumor suppressor APC2. Mol Cell Biochem, 2018, 443: 151–157

97 Guinot A, Oeztuerk-Winder F, Ventura J J. miR-17-92/p38α dysregulation enhances Wnt signaling and selects $Lgr6^+$ cancer stem-like cells during lung adenocarcinoma progression. Cancer Res, 2016, 76: 4012–4022

98 Zhu J, Wang S, Chen Y. miR-19 targeting of GSK3β mediates sulforaphane suppression of lung cancer stem cells. J Nutr Biochem, 2017, 44: 80–91

99 Yang F, Xiong H, Duan L, et al. MiR-1246 promotes metastasis and invasion of A549 cells by targeting GSK-3β–mediated Wnt/β-catenin pathway. Cancer Res Treat, 2019, 51: 1420–1429

100 Ren T, Fan X, Wang M, et al. miR-20b promotes growth of non−small cell lung cancer through a positive feedback loop of the Wnt/β−catenin signaling pathway. Int J Oncol, 2019, 56: 470–479

101 Zhu M C, Zhang Y H, Xiong P, et al. Circ-GSK3B up-regulates GSK3B to suppress the progression of lung adenocarcinoma. Cancer Gene Ther, 2022, 29: 1761–1772

102 Zhang H, Wang Y, Lu J, et al. Long non-coding RNA LINC00222 regulates GSK3β activity and promotes cell apoptosis in lung adenocarcinoma. Biomed Pharmacother, 2018, 106: 755–762

103 Koelman E M R, Yeste-Vázquez A, Grossmann T N. Targeting the interaction of β-catenin and TCF/LEF transcription factors to inhibit oncogenic Wnt signaling. Bioorg Med Chem, 2022, 70: 116920

104 Han X, Jiang H, Qi J, et al. Novel lncRNA UPLA1 mediates tumorigenesis and prognosis in lung adenocarcinoma. Cell Death Dis, 2020, 11: 999

105 Pei Y, He Y, Hu L, et al. The crosstalk between lncRNA-SNHG7/miRNA-181/cbx7 modulates malignant character in lung adenocarcinoma. Am J Pathol, 2020, 190: 1343–1354

106 Hu X, Zhou J, Chen X, et al. Pathway analyses identify novel variants in the WNT signaling pathway associated with tuberculosis in chinese population. Sci Rep, 2016, 6: 28530

107 Song Z, Wang H, Zhang S. Negative regulators of Wnt signaling in non-small cell lung cancer: theoretical basis and therapeutic potency. Biomed Pharmacother, 2019, 118: 109336

108 Jiang N, Zou C, Zhu Y, et al. HIF-1α-regulated miR-1275 maintains stem cell-like phenotypes and promotes the progression of LUAD by simultaneously activating Wnt/β-catenin and Notch signaling. Theranostics, 2020, 10: 2553–2570

109 Fang L, Cai J, Chen B, et al. Aberrantly expressed miR-582-3p maintains lung cancer stem cell-like traits by activating Wnt/β-catenin signalling. Nat Commun, 2015, 6: 8640

110 Zhang Y, Yuan J, Guo M, et al. Upregulation of long intergenic non-coding RNA LINC00326 inhibits non-small cell lung carcinoma progression by blocking Wnt/β-catenin pathway through modulating the miR-657/dickkopf WNT signaling pathway inhibitor 2 axis. Biol Direct, 2023, 18: 3

111 Li X, Lv F, Li F, et al. LINC01089 inhibits tumorigenesis and epithelial–mesenchymal transition of non-small cell lung cancer via the miR-27a/SFRP1/Wnt/β-catenin Axis. Front Oncol, 2020, 10: 532581

112 Yao Y, Hua Q, Zhou Y. CircRNA has_circ_0006427 suppresses the progression of lung adenocarcinoma by regulating miR-6783–3p/DKK1 axis and inactivating Wnt/β-catenin signaling pathway. Biochem Biophys Res Commun, 2019, 508: 37–45

113 Du X, Luo W, Li H, et al. Hsa_circ_0125356 promotes gemcitabine resistance by modulating WNT canonical and non-canonical pathways via miR-582-5p/FGF9 axis in non-small cell lung cancer. Mol Cancer, 2025, 24: 59

114 Zhang Q, Zhang B, Sun L, et al. MicroRNA-130b targets PTEN to induce resistance to cisplatin in lung cancer cells by activating Wnt/β-catenin pathway. Cell Biochem Function, 2018, 36: 194–202

115 Shi S L, Zhang Z H. Long non-coding RNA SNHG1 contributes to cisplatin resistance in non-small cell lung cancer by regulating miR-140-5p/Wnt/β-catenin pathway. Neoplasma, 2019, 66: 756–765

116 Wu Y, Cheng K, Liang W, et al. lncRNA RPPH1 promotes non–small cell lung cancer progression through the miR–326/WNT2B axis. Oncol Lett, 2020, 20: 1

117 Jiang P, Xu H, Xu C, et al. NEAT1 contributes to the CSC-like traits of A549/CDDP cells via activating Wnt signaling pathway. Chem-Biol Interact, 2018, 296: 154–161

118 Liao J, Lin J, Lin D, et al. Down-regulation of miR-214 reverses erlotinib resistance in non-small-cell lung cancer through up-regulating LHX6 expression. Sci Rep, 2017, 7: 781

119 Wang Y, Xia H, Zhuang Z, et al. Axl-altered microRNAs regulate tumorigenicity and gefitinib resistance in lung cancer. Cell Death Dis, 2014, 5: e1227

120 Rajtmajerová M, Trailin A, Liška V, et al. Long non-coding RNA and microRNA interplay in colorectal cancer and their effect on the tumor microenvironment. Cancers, 2022, 14: 5450

121 Alghazali T, Ahmed A T, Hussein U A R, et al. Noncoding RNA (ncRNA)-mediated regulation of TLRs: critical regulator of inflammation in tumor microenvironment. Med Oncol, 2025, 42: 144

122 Cen W, Yan Q, Zhou W, et al. miR-4739 promotes epithelial-mesenchymal transition and angiogenesis in “driver gene-negative” non-small cell lung cancer via activating the Wnt/β-catenin signaling. Cell Oncol, 2023, 46: 1821–1835

123 Luo H, Hu B, Gu X R, et al. The miR-23a/27a/24 – 2 cluster drives immune evasion and resistance to PD-1/PD-L1 blockade in non-small cell lung cancer. Mol Cancer, 2024, 23: 285

124 Taheriazam A, Bayanzadeh S D, Heydari Farahani M, et al. Non-coding RNA-based therapeutics in cancer therapy: an emphasis on Wnt/β-catenin control. Eur J Pharmacol, 2023, 951: 175781

125 Stewart J M. RNA nanotechnology on the horizon: self-assembly, chemical modifications, and functional applications. Curr Opin Chem Biol, 2024, 81: 102479

126 Nagatsugi F, Onizuka K. Selective chemical modification to the higher-order structures of nucleic acids. Chem Record, 2023, 23: e202200194

127 Anon. Modulating RNA with chemical modifications. Acc Chem Res, 2024, 57: 175–176

128 Rozners E. Chemical modifications of CRISPR RNAs to improve gene-editing activity and specificity. J Am Chem Soc, 2022, 144: 12584–12594
129 Li S, Li X, Xue W, et al. Screening for functional circular RNAs using the CRISPR–Cas13 system. Nat Methods, 2021, 18: 51–59
130 Yang H, Patel D J. Structures, mechanisms and applications of RNA-centric CRISPR–Cas13. Nat Chem Biol, 2024, 20: 673–688
131 Luo Y, Cook A B, Abdelmohsen L K E A, et al. Polymer vesicles and lipid nanoparticles. Annu Rev Mater Res, 2024, 54: 75–96
132 Yao Z, Liu T, Wang J, et al. Targeted delivery systems of siRNA based on ionizable lipid nanoparticles and cationic polymer vectors. Biotechnol Adv, 2025, 81: 108546
133 Huang X, Ding Y, Gu J, et al. Organ-selective lipid nanoparticles for precise cancer therapy: beyond liposomes and polymeric micelles. Chem Eng J, 2024, 494: 153171
134 Davodabadi F, Mirinejad S, Malik S, et al. Nanotherapeutic approaches for delivery of long non-coding RNAs: an updated review with emphasis on cancer. Nanoscale, 2024, 16: 3881–3914
135 Han S, Chen X, Huang L. The tumor therapeutic potential of long non-coding RNA delivery and targeting. Acta Pharm Sin B, 2023, 13: 1371–1382
136 Cina C, Majeti B, O'brien Z, et al. A novel lipid nanoparticle NBF-006 encapsulating glutathione S-transferase P siRNA for the treatment of KRAS-driven non–small cell lung cancer. Mol Cancer Ther, 2025, 24: 7–17
137 Viteri S, Rosell R. An innovative mesothelioma treatment based on miR-16 mimic loaded EGFR targeted minicells (TargomiRs). Transl Lung Cancer Res, 2018, 7: S1–S4
138 Payne D. RNA therapies. Nature, 2019, 574: S1
139 Winkle M, El-Daly S M, Fabbri M, et al. Noncoding RNA therapeutics — challenges and potential solutions. Nat Rev Drug Discov, 2021, 20: 629–651
140 Zhang P, Zhang D, Zhou W, et al. Network pharmacology: towards the artificial intelligence-based precision traditional Chinese medicine. Brief BioInf, 2023, 25: bbad518
141 Liao Y, Gui Y, Li Q, et al. The signaling pathways and targets of natural products from traditional Chinese medicine treating gastric cancer provide new candidate therapeutic strategies. Biochim Biophys Acta Rev Cancer, 2023, 1878: 188998
142 Yang Z, Zhang Q, Yu L, et al. The signaling pathways and targets of traditional Chinese medicine and natural medicine in triple-negative breast cancer. J EthnoPharmacol, 2021, 264: 113249
143 He G, Huang X, Dong Y, et al. Preliminary investigation on the mechanism of baicalein regulating the effects of Nischarin on invasion and apoptosis of human breast cancer cells MCF-7 through Wnt3α/β-catenin pathway. Int Immunopharmacol, 2024, 143: 113262
144 Jia Y, Chen L, Guo S, et al. Baicalin induced colon cancer cells apoptosis through miR-217/DKK1-mediated inhibition of Wnt signaling pathway. Mol Biol Rep, 2019, 46: 1693–1700
145 Han S H, Han J H, Chun W J, et al. Nobiletin inhibits non-small-cell lung cancer by inactivating WNT/β-catenin signaling through downregulating miR-15-5p. Evid-Based Complement Alternat Med, 2021, 2021: 1–9
146 Shi C, Zheng Y, Pan F, et al. Gallic acid suppressed tumorigenesis by an lncRNA MALAT1-Wnt/β-catenin axis in hepatocellular carcinoma. Front Pharmacol, 2021, 12: 708967
147 He Y, Liu X, Huang M, et al. Oleanolic acid inhibits the migration and invasion of hepatocellular carcinoma cells by promoting microRNA-122 expression. Pharmazie, 2021, 76: 422–427
148 Su Z, Hu B, Li J, et al. Paeoniflorin inhibits colorectal cancer cell stemness through the miR-3194-5p/catenin beta-interacting protein 1 axis. Kaohsiung J Med Scie, 2023, 39: 1011–1021
149 Jing W, Dong H, Min M, et al. Dependence of artesunate on long noncoding RNA-RP11 to inhibit epithelial-mesenchymal transition of hepatocellular carcinoma. J Cell Biochem, 2019, 120: 6026–6034
150 Zhu D, Li A, Lv Y, et al. Traditional chinese medicine: a class of potentially reliable epigenetic drugs. Front Pharmacol, 2022, 13: 907031
151 Zhang Z J, Song X G, Xie L, et al. Circulating serum exosomal miR-20b-5p and miR-3187-5p as efficient diagnostic biomarkers for early-stage non-small cell lung cancer. Exp Biol Med (Maywood), 2020, 245: 1428–1436
152 Aval S F, Lotfi H, Sheervalilou R, et al. Tuning of major signaling networks (TGF-β, Wnt, Notch and Hedgehog) by miRNAs in human stem cells commitment to different lineages: possible clinical application. Biomed Pharmacother, 2017, 91: 849–860
153 Sonawala K, Ramalingam S, Sellamuthu I. Influence of long non-coding RNA in the regulation of cancer stem cell signaling pathways. Cells, 2022, 11: 3492

Summary for "非小细胞肺癌Wnt/β-catenin信号通路相关ncRNAs: 多维机制解析与诊疗转化研究"

NcRNAs in the Wnt/β-catenin signaling pathway: from multidimensional mechanisms to diagnostic and therapeutic translation in non-small-cell lung cancer

Yang Zhong[1], Heng-Zhou Lai[1], Jia-Wei He[1], Chun-Xia Huang[1], Ao-Ling Yang[1], Wen-Bo Huang[1], Qiong Ma[1,3], Xi Fu[1,2,3], Chuan Zheng[1*] & Feng-Ming You[1,2,3*]

[1] *TCM Prevention and Treatment of Metabolic and Chronic Diseases Key Laboratory of Sichuan Province, Hospital of Chengdu University of Traditional Chinese Medicine, Chengdu 610075, China*
[2] *Oncology Teaching and Research Department, Chengdu University of Traditional Chinese Medicine, Chengdu 610075, China*
[3] *Institute of Oncology, Chengdu University of Traditional Chinese Medicine, Chengdu 610075, China*
* Corresponding authors, E-mail: zhengchuan@cdutcm.edu.cn; yfmdoc@163.com

Lung cancer (LC) is the leading cause of cancer-related deaths worldwide, with non-small-cell lung cancer (NSCLC) constituting the majority of cases. Effective treatment is hampered by an incomplete understanding of molecular mechanisms, insufficient early-detection tools, and widespread drug resistance. The Wnt/β-catenin signaling pathway, central to lung tissue homeostasis, is frequently dysregulated in NSCLC and drives malignant transformation and progression. Accumulating evidence indicates that non-coding RNAs (ncRNAs)—including microRNAs (miRNAs), long non-coding RNAs (lncRNAs), and circular RNAs (circRNAs)—are critical modulators of this pathway. However, the multilayered regulatory networks they form, as well as strategies for their clinical translation, remain incompletely defined.

This review synthesizes current knowledge on how diverse ncRNAs target the Wnt/β-catenin pathway in NSCLC, addressing their impacts on ligand secretion, receptor activation, intracellular signal transduction, and negative regulators. It outlines a multidimensional regulatory landscape in which ncRNAs collaboratively promote or suppress NSCLC by modulating Wnt/β-catenin signaling and evaluates their potential as biomarkers and therapeutic targets.

Mechanistically, ncRNAs operate at transcriptional, post-transcriptional, translational, and epigenetic levels to influence cancer stem cell maintenance, epithelial–mesenchymal transition, angiogenesis, proliferation, metastasis, and therapy resistance. miRNAs commonly bind the 3′ untranslated regions of mRNAs encoding pathway components–such as secreted frizzled-related proteins, Dickkopf family members, GSK-3β, AXIN1, and β-catenin–thereby fine-tuning their expression and directly influencing malignant behaviors. Many lncRNAs act as competing endogenous RNAs (ceRNAs), sequestering specific miRNAs to relieve repression of target transcripts and activate Wnt signaling; others function as signals, scaffolds, decoys, or guides to regulate transcription factors and pathway components. Owing to their covalently closed structure, circRNAs are stable and primarily function as miRNA sponges; a subset also encodes functional peptides that influence phosphorylation, ubiquitination, or stability of Wnt/β-catenin components.

Therapeutically, ncRNAs function as pivotal regulatory nodes and represent promising targets for intervention. They are centrally involved in mechanisms of drug resistance, indicating that targeting the ncRNA–Wnt/β-catenin axis could synergize with existing treatments. Advances in RNA delivery platforms and RNA-based therapeutics offer promising routes to reverse resistance and remodel the tumor microenvironment. Additionally, natural compounds with multi-target activity and small molecules that modulate specific ncRNAs provide complementary strategies to accelerate drug discovery and personalize therapy.

In diagnostics, Wnt/β-catenin-related ncRNAs show considerable promise as clinical biomarkers throughout the management of NSCLC, including early detection, molecular subtyping, metastasis surveillance, recurrence monitoring, prognosis, and resistance assessment. High-throughput sequencing and comprehensive expression profiling have identified candidate ncRNA biomarkers showing promise for high sensitivity and specificity in preliminary studies, which require rigorous clinical validation.

Integrated multi-omics and functional studies are needed to decode ncRNA regulatory networks in NSCLC systematically. Establishing a translational pipeline from "mechanistic discovery → targeted therapeutic development → biomarker validation" will be critical to moving ncRNA findings from bench to bedside and ultimately improving the diagnosis and treatment of NSCLC.

Wnt/β-catenin, ncRNAs, non-small-cell lung cancer (NSCLC), mechanism of action, biomarkers

doi: 10.1360/CSB-2025-0706