SCIENCE CHINA
Chemistry

• ARTICLES •

July 2013 Vol.56 No.7: 945–951
doi: 10.1007/s11426-013-4843-7

Synthesis of naphthalene derivatives through inexpensive BF_3·Et_2O-catalyzed annulation reaction of arylacetaldehydes with arylalkynes

XIANG ShiKai[1*], HU Hao[1], MA Jing[1], LI YuanZhuo[1], WANG BiQin[1*], FENG Chun[1], ZHAO KeQing[1], HU Ping[1] & CHEN XiaoZhen[2]

[1]*College of Chemistry and Materials Science, Sichuan Normal University, Chengdu 610066, China*
[2]*Chengdu Institute of Biology, Chinese Academy of Sciences, Chengdu 610041, China*

Received November 11, 2012; accepted December 26, 2012; published online February 21, 2013

An inexpensive BF_3·Et_2O-catalyzed annulation reaction of arylacetaldehydes with arylalkynes has been developed. Various substituted phenylacetaldehydes and phenylacetylenes can undergo this reaction, producing corresponding α-aryl substituted naphthalene derivatives. Use of inexpensive and readily available BF_3·Et_2O catalyst constitutes the most attractive advantage of this transformation.

naphthalene derivatives, BF_3·Et_2O, arylacetaldehydes, arylalkynes, annulation reaction


1 Introduction

The naphthalene unit is a ubiquitous skeleton in chemical and pharmaceutical industries as well as optical and electronic materials [1]. In the past decades, the development of some new and efficient methodologies for the synthesis of polysubstituted naphthalene derivatives has aroused great interest of organic chemists [2–5]. Among these methodologies, the acid-catalyzed annulation reaction of α-aryl-substituted carbonyl compounds with alkynes is quite an efficient approach for the synthesis of naphthalene derivatives. In 2002, Li and co-workers [3] realized a highly regioselective synthesis of polysubstituted naphthalene derivatives through $GaCl_3$-catalyzed annulation reaction of phenylacetaldehydes (and ketones) with alkynes (Eq. (1)). In spite of the moderate yields and the application of expensive $GaCl_3$ catalyst, this annulation reaction was discovered by them for the first time. Next year, Kabalka and co-workers [4] reported a $TiCl_4$-mediated annulation reaction of α-aryl-substituted carbonyl compounds with alkynes. Kabalka's method greatly improved the yields of naphthalene derivatives, but required the use of quantitative $TiCl_4$ for this transformation. In 2009, the catalytic system of $AuCl_3/AgSbF_6$ was utilized by Balamurugan group to achieve the similar transformation to synthesize polysubstituted naphthalene derivatives (Eq. (2)) [5]. This approach is very effective, however, use of expensive $AuCl_3/AgSbF_6$ catalyst limited its broad application.

Inexpensive BF_3·Et_2O is widely used as a Lewis acid catalyst in organic synthesis [6]. Recently, we developed a BF_3·Et_2O/ammonium salt cocatalyzed alkenylation reaction of indoles with α,β-unsaturated ketones [7]. The combination of inexpensive and readily available BF_3·Et_2O and an ammonium salt as the efficient cocatalyst constitutes the attractive advantage of the reaction. Herein, we wish to report a new annulation reaction of arylacetaldehydes with arylalkynes catalyzed by inexpensive BF_3·Et_2O to give naphthalene derivatives (Eq. (3)).

*Corresponding authors (email: xiangsk@hotmail.com; wangbiqin1964@126.com)


© Science China Press and Springer-Verlag Berlin Heidelberg 2013



2 Results and discussion

Initially, we investigated the annulation reaction of phenylacetaldehyde **1a** and phenylacetylene **2a** in the presence of 5 mol% of $FeCl_3$. Gratifyingly, 58% of 1-phenylnaphthalene **3aa** was obtained when dichloroethane (DCE) was used as the solvent (entry 1, Table 1). Although other Lewis acid catalysts or Brønsted acid catalysts such as $FeCl_2$·$4H_2O$, AgOTf, and CH_3SO_3H also promoted this transformation, their efficiencies were lower than $FeCl_3$ (entries 2–4, Table 1). TsOH hardly led to any desired product (entry 5, Table 1). Notably, the yield was improved to 74% when BF_3·Et_2O was used as a catalyst (entry 6, Table 1).

Table 1 Annulation reaction of phenylacetaldehyde (**1a**) with phenylacetylene (**2a**) under different conditions [a)]




Entry	Catalyst (mmol%)	Solvent (mL)	*T* (℃)	Yield (%) [b)]
1	$FeCl_3$ (5)	DCE	80	58
2	$FeCl_2 \cdot H_2O$ (5)	DCE	80	27
3	AgOTf (5)	DCE	80	27
4	CH_3SO_3H (5)	DCE	80	28
5	TsOH (5)	DCE	80	trace
6	$BF_3 \cdot Et_2O$ (5)	DCE	80	74
7	$BF_3 \cdot Et_2O$ (5)	PhCl	80	42
8	$BF_3 \cdot Et_2O$ (5)	toluene	80	26
9	$BF_3 \cdot Et_2O$ (5)	MeCN	80	38
10	$BF_3 \cdot Et_2O$ (5)	$MeNO_2$	80	37
11	$BF_3 \cdot Et_2O$ (5)	DMF	80	ND [c)]
12	$BF_3 \cdot Et_2O$ (5)	DCE	50	30
13	$BF_3 \cdot Et_2O$ (5)	DCE	rt	ND [c)]
14[d]	$BF_3 \cdot Et_2O$ (5)	DCE	80	82













a) All the reactions were carried out in the scale of 0.3 mmol **1a**, 0.3 mmol **2a**, 5 mmol% catalyst in 1.0 mL solvent if without further note; b) the yields were isolated yields; c) ND = not detected; d) 0.36 mmol **2a** was used.

Different solvents were then screened, indicating DCE as the best reaction medium (entries 6–11, Table 1). In addition, the reaction temperature was also critical for this annulation reaction (cf. entries 6, 12 and 13, Table 1). Finally, the best result of 82% yield was achieved by increasing the amount of phenylacetylene to 1.2 equivalents (entry 14, Table 1).

The scope of annulation reaction was expanded to a variety of substituted phenylacetaldehydes **1** and phenylacetylenes **2** (Tables 2 and 3). Phenylacetaldehydes **1** with both electron-donating groups and electron-withdrawing groups in the benzene ring smoothly underwent this kind of transformation, generating naphthalene derivatives **3** in moderate


Table 2 Annulation reaction of substituted phenylacetaldehydes (**1**) with phenylacetylene (**2a**) [a)]



entry	1		3		yield (%)
1		**1a**		**3aa**	82
2		**1b**		**3ba**	58
3		**1c**		**3ca**	60
4[b]		**1d**		**3da**	30
5		**1e**		**3ea**	53
6		**1f**		**3fa**	64

a) All the reactions were carried out in the scale of 0.3 mmol **1**, 0.36 mmol **2a**, 5 mmol% $BF_3 \cdot Et_2O$ in 1.0 mL DCE at 80 °C if without further note. The yields were isolated yields; b) the reactions were carried out at 90 °C. 10 mmol% $BF_3 \cdot Et_2O$ was used.

Table 3 Annulation reaction of phenylacetaldehydes (**1a**) with various phenylacetylenes (**2**) [a)]




1a 2 3

entry	2	3	yield (%)
1	**2a**	**3aa**	82
2	**2b**	**3ab**	73
3	**2c**	**3ac**	78
4 [b)]	**2d**	**3ad**	55
5 [b)]	**2e**	**3ae**	61
6	**2f**	**3af**	0
7	**2g**	**3ag**	0
8	**2h**	**3ah**	trace

a) All the reactions were carried out in the scale of 0.3 mmol **1a**, 0.36 mmol **2**, 5 mmol% $BF_3 \cdot Et_2O$ in 1.0 mL DCE at 80 °C if without further note. The yields were isolated yields; b) the reactions were carried out in PhCl (1 mL) instead of DCE at 110 °C. 20 mmol% $BF_3 \cdot Et_2O$ was used.


to excellent yields 30–82% (entries 1–6, Table 2). It is noteworthy that electronic effect of the arylacetaldehydes component had some influence on the reaction, but the regularity was not obvious (entries 1–6, Table 2). The substituents Cl and Br were compatible under these conditions, which could be further transformed into other functionalities (entries 5 and 6, Table 2).

In addition, both electron-rich and electron-deficient arylacetylenes were tolerable as the annulation reaction partners with the yields of 55–82% (entries 1–5, Table 3). Notably, the electron-rich arylacetylenes gave the desired products in higher yields than the electron-deficient ones (entries 2–5, Table 3). It is regretful that both (trimethylsilyl)acetylene and internal alkyne can not undergo this transformation to give the desired product (entries 6 and 7, Table 3). The internal alkynes failed to undergo the reaction possibly due to the following two reasons: i) The diphenylacetylene has a weaker nucleophilicity than phenylacetaldehyde; ii) The Lewis acidity of $BF_3 \cdot Et_2O$ is lower than $GaCl_3$ or $AuCl_3/AgSbF_6$. Furthermore, the 1-hexyne **2h** was examined and gave only trace amount of product (entry 8, Table 3).



Two meta-substituted arylacetaldehydes (**1g**, **1h**) was employed to study the selectivity of the reaction (Eq. (4)). The results showed that the selectivity is terrible.


To explore the application of such transformations, this annulation reaction was scaled up to 10 mmol (Eq. (5)). When 10 mmol of phenylacetaldehyde **1a** and phenylacetylene **2a** were used as substrates, 1.32 g of 1-phenylnaphthalene **3aa** was obtained with the good yield of 65%. The result indicated that the catalytic method was very efficient.



A possible mechanism for this catalytic transformation is illustrated in Scheme 1. Initially, $BF_3 \cdot Et_2O$ coordinates with the carbonyl oxygen to form the aldehyde-Lewis acid complex **A** [3, 5, 6]. Electrophilic attack of the carbonyl carbon in complex **A** on the arylalkyne **2** takes place to generate complex **B** with a vinyl carbocation stabilized by the aryl group. Intramolecular electrophilic attack of the formed vinyl carbocation to the aromatic ring followed by elimination of a proton generates the cyclization product **D**. Finally, the complex **D** can aromatize by dehydration to form the corresponding naphthalene derivative **3**, releasing the catalyst at the completion of catalytic cycle.

3 Conclusions

In summary, we developed an inexpensive $BF_3 \cdot Et_2O$-catalyzed annulation reaction of arylacetaldehydes with arylalkynes.



Scheme 1 Proposed mechanism for this catalytic transformation.

Various substituted phenylacetaldehydes and phenylacetylenes can undergo this reaction, providing an alternative approach for the synthesis of naphthalene derivatives. Use of inexpensive and readily available BF_3·Et_2O catalyst constitutes the most attractive advantage of this transformation. Development of other methodologies for the synthesis of naphthalene derivatives is ongoing in our laboratory.


4 Experimental

4.1 General experimental section

All manipulations were conducted with Schlenk tube. ^{1}H NMR spectra were recorded on the Varian 400 MHz WB spectrometers. Chemical shifts (in ppm) were referenced to tetramethylsilane (δ = 0 ppm) in $CDCl_3$ as an internal standard. ^{13}C NMR spectra were obtained by the same NMR spectrometers and were calibrated with $CDCl_3$ (δ = 77.00 ppm). Mass spectra were obtained using Electron Impact (EI) mass spectrometer. Arylacetaldehydes **1b–f** were synthesized according to literature method [8]. Unless otherwise noted, materials and solvents from commercial suppliers were used without further purification.


4.2 Experimental procedures and characterization of products

*1-Phenylnaphthalene (**3aa**)*




Typical procedure: phenylacetaldehyde (36.0 mg, 0.3 mmol) and phenylacetylene (36.7 mg, 0.36 mmol), BF_3·Et_2O (2.1 mg, 0.015 mmol) and DCE (1 mL) were mixed in a Schlenck tube. The reaction mixture was stirred for 15 h at 80 °C. The solution was cooled to room temperature and the solvent was removed under vacuum. The crude product was purified by column chromatography on silica gel (eluent: petroleum ether/dichloromethane, *v*/*v* = 70:1) to afford 50.2 mg (82%) of **3aa**. IR:(KBr) ν_{max} 3056, 2921, 2851 cm^{-1}; ^{1}H NMR (400 MHz, $CDCl_3$, ppm) δ 7.91–7.88 (m, 2H), 7.85–7.82 (m, 1H), 7.48–7.40 (m, 9H); ^{13}C NMR (100 MHz, $CDCl_3$, ppm) δ 140.7, 140.2, 133.7, 131.6, 130.0, 128.2, 127.6, 127.2, 126.9, 126.0, 125.7, 125.4; MS (70 eV), *m*/*z* (%): 204.4 (72) $[M]^+$, 203.2 (100).

*7-Methyl-1-phenylnaphthalene (**3ba**)*



2-*p*-Tolylacetaldehyde (40.2 mg, 0.3 mmol) and phenylacetylene (36.7 mg, 0.36 mmol), BF_3·Et_2O (2.1 mg, 0.015 mmol) and DCE (1 mL) were mixed in a Schlenck tube. The reaction mixture was stirred for 15 h at 80 °C. The solution was cooled to room temperature and the solvent was removed under vacuum. The crude product was purified by column chromatography on silica gel (eluent: petroleum ether/dichloromethane, *v*/*v* = 70:1) to afford 37.9 mg (58%) of **3ba**. IR(KBr): ν_{max} 3052, 2917, 2855, 2732 cm^{-1}; ^{1}H NMR (400 MHz, $CDCl_3$, ppm) δ 7.79 (d, *J* = 8.0 Hz, 2H), 7.66 (s, 1H), 7.50–7.45 (m, 4H), 7.44–7.41 (m, 2H), 7.38–7.36 (m, 1H), 7.33–7.30 (m, 1H), 2.42 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$, ppm) δ 141.0, 139.5, 135.7, 132.0, 131.7, 130.0, 128.2, 128.1, 128.0, 127.4, 127.1, 127.0, 124.8, 124.5, 21.9; MS (70 eV), *m*/*z* (%): 218.0 (100) $[M]^+$.


*7-Methoxy-1-phenylnaphthalene (**3ca**)*



2-(4-Methoxyphenyl)acetaldehyde (45.0 mg, 0.3 mmol) and phenylacetylene (36.7 mg, 0.36 mmol), BF_3·Et_2O (2.1 mg, 0.015 mmol) and DCE (1 mL) were mixed in a Schlenck tube. The reaction mixture was stirred for 15 h at 80 °C. The solution was cooled to room temperature and the solvent was removed under vacuum. The crude product was purified by column chromatography on silica gel (eluent: petroleum ether/dichloromethane, *v*/*v* = 5:1) to afford 42.1 mg (60%) of **3ca** with two isomers in 92:8 ratio. IR(KBr) ν_{max} 3054, 3022, 2956, 2936, 2878, 2831 cm^{-1}; ^{1}H NMR of major product (400 MHz, $CDCl_3$, ppm) δ 7.81–7.79 (m, 2H), 7.52–7.46 (m, 4H), 7.43–7.41 (m, 1H), 7.39–7.36 (m, 2H),


7.23 (s, 1H), 7.16 (d, J = 8.8 Hz, 1H), 3.75 (s, 3H); MS (70 eV), *m*/*z* (%): 234.1 (100) $[M]^+$.

*1-Methoxy-5-phenylnaphthalene (**3da**)*




2-(2-Methoxyphenyl)acetaldehyde (45.0 mg, 0.3 mmol) and phenylacetylene (36.7 mg, 0.36 mmol), BF_3·Et_2O (4.2 mg, 0.03 mmol) and DCE (1 mL) were mixed in a Schlenck tube. The reaction mixture was stirred for 15 h at 90 °C. The solution was cooled to room temperature and the solvent was removed under vacuum. The crude product was purified by column chromatography on silica gel (eluent: petroleum ether/dichloromethane, *v*/*v* = 5:1) to afford 21.1 mg (30%) of **3da**. IR(KBr) ν_{max} 3031, 3006, 2966, 2934, 2853, 2833 cm^{-1}; ^1H NMR (400 MHz, $CDCl_3$, ppm) δ 8.33–8.29 (m, 1H), 7.53–7.41 (m, 8H), 7.33–7.28 (m, 1H), 6.81 (d, J = 7.2 Hz, 1H), 3.99 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$, ppm) δ 155.5, 141.1, 139.8, 132.6, 130.1, 128.1, 127.5, 127.1, 125.9, 125.8, 124.7, 121.5, 118.3, 103.6, 55.6; MS (70 eV), *m*/*z* (%): 234.2 (100) $[M]^+$.

*7-Chloro-1-phenylnaphthalene (**3ea**)*




2-(4-Chlorophenyl)acetaldehyde (46.4 mg, 0.3 mmol) and phenylacetylene (36.7 mg, 0.36 mmol), BF_3·Et_2O (2.1 mg, 0.015 mmol) and DCE (1 mL) were mixed in a Schlenck tube. The reaction mixture was stirred for 15 h at 80 °C. The solution was cooled to room temperature and the solvent was removed under vacuum. The crude product was purified by column chromatography on silica gel (eluent: petroleum ether/dichloromethane, *v*/*v* = 70:1) to afford 41.6 mg (58%) of **3ea**. IR(KBr) ν_{max} 3056, 2924, 2853 cm^{-1}; ^1H NMR (400 MHz, $CDCl_3$, ppm) δ 7.87–7.81 (m, 3H), 7.51–7.44 (m, 8H); ^{13}C NMR (100 MHz, $CDCl_3$, ppm) δ 140.0, 139.6, 132.3, 132.01, 131.99, 129.93, 129.86, 128.4, 127.9, 127.5, 127.4, 126.7, 125.6, 124.9; MS (70 eV), *m*/*z* (%): 238.0 (100) $[M]^+$.

*7-Bromo-1-phenylnaphthalene (**3fa**)*




2-(4-Bromophenyl)acetaldehyde (59.7 mg, 0.3 mmol) and phenylacetylene (36.7 mg, 0.36 mmol), BF_3·Et_2O (2.1 mg, 0.015 mmol) and DCE (1 mL) were mixed in a Schlenck tube. The reaction mixture was stirred for 15 h at 80 °C. The solution was cooled to room temperature and the solvent was removed under vacuum. The crude product was purified by column chromatography on silica gel (eluent: petroleum ether/dichloromethane, *v*/*v* = 70:1) to afford 54.0 mg (64%) of **3fa**. IR(KBr) ν_{max} 3056, 2922, 2851 cm^{-1}; ^1H NMR (400 MHz, $CDCl_3$, ppm) δ 8.04 (s, 1H), 7.80 (d, J = 8.0 Hz, 1H), 7.75 (d, J = 8.4 Hz, 1H), 7.57–7.43 (m, 8H); ^{13}C NMR (100 MHz, $CDCl_3$, ppm) δ 140.0, 139.5, 132.7, 132.2, 129.9, 129.2, 128.5, 128.1, 127.9, 127.53, 127.48, 125.8, 120.4; MS (70 eV), *m*/*z* (%): 282.0 (100) $[M]^+$.

*1-p-Tolylnaphthalene (**3ab**)*



Phenylacetaldehyde (36.0 mg, 0.3 mmol) and 1-ethynyl-4-methylbenzene (41.8 mg, 0.36 mmol), BF_3·Et_2O (2.1 mg, 0.015 mmol) and DCE (1 mL) were mixed in a Schlenck tube. The reaction mixture was stirred for 15 h at 80 °C. The solution was cooled to room temperature and the solvent was removed under vacuum. The crude product was purified by column chromatography on silica gel (eluent: petroleum ether/dichloromethane, *v*/*v* = 70:1) to afford 47.7 mg (73%) of **3ab**. IR(KBr) ν_{max} 3037, 2963, 2918, 2857 cm^{-1}; ^1H NMR (400 MHz, $CDCl_3$): δ 7.92–7.85 (m, 3H), 7.51–7.41 (m, 6H), 7.31 (s, 2H), 2.46 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$): δ 140.2, 137.8, 136.9, 133.8, 131.7, 130.0, 129.0, 128.3, 127.4, 126.9, 126.1, 125.9, 125.7, 125.4, 21.3; MS (70 eV), *m*/*z* (%): 218.2 (100) $[M]^+$.


*1-(4-Tert-butylphenyl)naphthalene (**3ac**)*



Phenylacetaldehyde (36.0 mg, 0.3 mmol) and 1-*tert*-butyl-4-ethynylbenzene (57.0 mg, 0.36 mmol), BF_3·Et_2O (2.1 mg, 0.015 mmol) and DCE (1 mL) were mixed in a Schlenck tube. The reaction mixture was stirred for 15 h at 80 °C. The solution was cooled to room temperature and the solvent was removed under vacuum. The crude product was purified by column chromatography on silica gel (eluent: petroleum ether/dichloromethane, *v*/*v* = 70:1) to afford 60.9 mg (78%) of **3ac**. IR(KBr) ν_{max} 3063, 3039, 2958, 2900, 2863 cm^{-1}; ^1H NMR (400 MHz, $CDCl_3$, ppm) δ 7.96 (d, J = 8.8 Hz, 1H), 7.90 (d, J = 8.4 Hz, 1H), 7.84 (d, J = 8.0 Hz, 1H), 7.53–7.48 (m, 4H), 7.47–7.40 (m, 4H), 1.41 (s, 9H); ^{13}C NMR (100 MHz, $CDCl_3$, ppm) δ 150.1, 140.2, 137.7, 133.8,


131.7, 129.7, 128.2, 127.4, 126.9, 126.2, 125.9, 125.7, 125.4, 125.2, 34.6, 31.5; MS (70 eV), m/z (%): 260.0 (100) $[M]^+$.

*1-(4-Bromophenyl)naphthalene (**3ad**)*




Phenylacetaldehyde (36.0 mg, 0.3 mmol) and 1-bromo-4-ethynylbenzene (65.2 mg, 0.36 mmol), $BF_3{\cdot}Et_2O$ (8.2 mg, 0.06 mmol) and PhCl (1 mL) were mixed in a Schlenck tube. The reaction mixture was stirred for 15 h at 110 °C. The solution was cooled to room temperature and the solvent was removed under vacuum. The crude product was purified by column chromatography on silica gel (eluent: petroleum ether/dichloromethane, *v*/*v* = 70:1) to afford 46.7 mg (55%) of **3ad**. IR(KBr) ν_{max} 3058, 2924 cm^{-1}; ^1H NMR (400 MHz, $CDCl_3$, ppm) δ 7.90–7.81 (m, 3H), 7.61–7.58 (m, 2H), 7.49–7.47 (m, 2H), 7.44–7.41 (m, 1H), 7.35–7.33 (m, 3H); ^{13}C NMR (100 MHz, $CDCl_3$, ppm) δ 139.6, 138.9, 133.7, 131.7, 131.4, 131.3, 128.3, 128.0, 126.8, 126.2, 125.9, 125.6, 125.3, 121.4; MS (70 eV), *m*/*z* (%): 282.2 (100) $[M]^+$.

*1-(4-Fluorophenyl)naphthalene (**3ae**)*




Phenylacetaldehyde (36.0 mg, 0.3 mmol) and 1-ethynyl-4-fluorobenzene (43.2 mg, 0.36 mmol), $BF_3{\cdot}Et_2O$ (8.4 mg, 0.06 mmol) and PhCl (1 mL) were mixed in a Schlenck tube. The reaction mixture was stirred for 15 h at 110 °C. The solution was cooled to room temperature and the solvent was removed under vacuum. The crude product was purified by column chromatography on silica gel (eluent: petroleum ether/dichloromethane, *v*/*v* = 70:1) to afford 40.6 mg (61%) of **3ae**. IR(KBr) ν_{max} 3042, 2929, 2850 cm^{-1}; ^1H NMR (400 MHz, $CDCl_3$): δ 7.91–7.84 (m, 3H), 7.52–7.36 (m, 6H), 7.19–7.14 (m, 2H); ^{13}C NMR (100 MHz, $CDCl_3$): δ 162.2 (d, *J* = 244.4 Hz), 139.1, 136.63, 136.59, 133.7, 131.6, 131.5, 128.3, 127.8, 127.0, 126.1, 125.8, 125.7, 125.3, 115.3, 115.0; MS (70 eV), m/z (%): 222.4 (74) $[M]^+$, 221.1 (100).

*6-Chloro-1-phenylnaphthalene (**3ga-1**) and 1-chloro-8-phenylnaphthalene (**3ga-2**)*



2-(3-Chlorophenyl)acetaldehyde (46.4 mg, 0.3 mmol) and phenylacetylene (36.7 mg, 0.36 mmol), $BF_3{\cdot}Et_2O$ (2.1 mg, 0.015 mmol) and DCE (1 mL) were mixed in a Schlenck tube. The reaction mixture was stirred for 15 h at 80 °C. The solution was cooled to room temperature and the solvent was removed under vacuum. The crude product was purified by column chromatography on silica gel (eluent: petroleum ether/dichloromethane, *v*/*v* = 70:1) to afford 21.4 mg (30%) of **3ga** (**3ga-1/3ga-2 =** 61/39). **3ga-1:** ^1H NMR (400 MHz, $CDCl_3$, ppm) δ 7.867 (d, *J* = 2.0 Hz, 1H), 7.82 (d, *J* = 8.8 Hz, 1H), 7.75 (d, *J* = 8.0 Hz, 1H), 7.55–7.39 (m, 7H), 7.34 (dd, J_1 = 9.6 Hz, J_2 = 2.0 Hz, 1H); ^{13}C NMR (100 MHz, $CDCl_3$, ppm) δ 140.4, 140.2, 134.5, 131.6, 129.93, 129.0, 128.3, 127.8, 127.5, 127.1, 126.8, 126.7, 126.5; MS (70 eV), *m*/*z* (%): 237.7 (100) $[M]^+$. **3ga-2:** ^1H NMR (400 MHz, $CDCl_3$, ppm) δ 7.88–7.82 (m, 2H), 7.52–7.47 (m, 2H), 7.41–7.32 (m, 7H); ^{13}C NMR (100 MHz, $CDCl_3$, ppm) δ 143.5, 139.5, 136.0, 131.4, 131.0, 129.6, 129.4, 128.7, 128.6, 128.2, 127.2, 126.7, 125.6, 125.4; MS (70 eV), *m*/*z* (%): 238.7 (100) $[M]^+$.

*6-Methyl-1-phenylnaphthalene (**3ha-1**) and 1-methyl-8-phenylnaphthalene (**3ha-2**)*



2-*m*-Tolylacetaldehyde (40.2 mg, 0.3 mmol) and phenylacetylene (36.7 mg, 0.36 mmol), $BF_3{\cdot}Et_2O$ (2.1 mg, 0.015 mmol) and DCE (1 mL) were mixed in a Schlenck tube. The reaction mixture was stirred for 15 h at 80 °C. The solution was cooled to room temperature and the solvent was removed under vacuum. The crude product was purified by column chromatography on silica gel (eluent: petroleum ether/dichloromethane, *v*/*v* = 70:1) to afford 28.8 mg (44%) of **3ha** (**3ha-1/3ha-2 =** 53/47). ^1H NMR (400 MHz, $CDCl_3$, ppm) δ 7.66 (s, 1H); MS (70 eV), *m*/*z* (%): 218.2 (100) $[M]^+$.

We are grateful for the financial supports from Sichuan Provincial Department of Education (11ZA108), National Natural Science Foundation of China (21202109 and 21072140), Special Funds of Sichuan Normal University for Sharing the Large Precision Equipments (DJ2012-07, DJ2011-16) and Sichuan Normal University.

1 (a) Medarde M, Maya ABS, Pérez-Melero CJ. Naphthalene combretastatin analogues: Synthesis, cytotoxicity and antitubulin activity. *Enzyme Inhib Med Chem*, 2004, 19: 521–540; (b) Xie X, Kozlowski MC. Synthesis of the naphthalene portion of the rubromycins. *Org Lett*, 2001, 3: 2661–2663; (c) Ukita T, Nakamura Y, Kubo A, Yamamoto Y, Takahashi M, Kotera J, Ikeo T. 1-Arylnaphthalene lignan: A novel scaffold for type 5 phosphodiesterase inhibitor. *J Med Chem*, 1999, 42: 1293–1305; (d) Ward RS. Lignans, neolignans, and related compounds. *Nat Prod Rep*, 1995, 12: 183–205; (e) Trujillo JM, Jorge RE, Navarro E, Boada J. Lignans from Justicia hyssopifolia. *Phytochemistry*, 1990, 29: 2991–2993; (f) Seong BL, Han

MH. A facile preparation of rifamycin derivatives by use of manganese dioxide. *Chem Lett*, 1982, 627–628; (g) Georghiou PE, Li Z, Ashram M, Chowdhury S, Mizyed S, Tran AH, Al-Saraierh H, Miller DO. Calixnaphthalenes: Deep, electron-rich naphthalene ring-containing calixarenes. the first decade. *Synlett*, 2005, 879–891; (h) Watson MD, Fechtenkötter A, Müllen K. Big is beautiful–"aromaticity" revisited from the viewpoint of macromolecular and supramolecular benzene chemistry. *Chem Rev*, 2001, 101: 1267–1300; (i) Zeng F, Su YS, Chen CF. Li+-templated complexation of cylindrical macrotricyclic host with naphthalene diimide: Cation-controlled switchable complexation processes. *Sci China Chem*, 2012, 10: 2069–2074

2 For some reviews, see: (a) Koning CB, Rousseau AL, Otterlo WAL. Modern methods for the synthesis of substituted naphthalenes. Tetrahedron, 2003, 59: 7–36; (b) Saito S, Yamamoto Y. Recent advances in the transition-metal-catalyzed regioselective approaches to polysubstituted benzene derivatives. *Chem Rev*, 2000, 100: 2901–2916; (c) Bradsher CK. Formation of six-membered aromatic rings by cyclialkylation of some aldehydes and ketones. *Chem Rev*, 1987, 87: 1277–1297; (d) Asao N. Gold- and copper-catalyzed [4+2] benzannulations between enynal or enynone units and 2π-systems. *synlett*, 2006, 1645–1656.

3 (a) Viswanathan GS, Wang M, Li CJ. A highly regioselective synthesis of polysubstituted naphthalene derivatives through Gallium trichloride catalyzed alkyne–aldehyde coupling. *Angew Chem Int Ed*, 2002, 41: 2138–2141; (b) Viswanathan GS, Li CJ. Synthesis of naphthalene derivatives via a novel Gallium trichloride catalyzed cross-coupling of epoxides with alkynes. *synlett*, 2002, 1553–1555

4 Kabalka GW, Ju Y, Wu Z. A new Titanium tetrachloride mediated annulation of α-aryl-substituted carbonyl compounds with alkynes: A simple and highly efficient method for the regioselective synthesis of polysubstituted naphthalene derivatives. *J Org Chem*, 2003, 68: 7915–7917

5 Balamurugan R, Gudla V. Gold-catalyzed electrophilic addition to arylalkynes. A facile method for the regioselective synthesis of substituted naphthalenes. *Org Lett*, 2009, 11: 3116–3119; For the intramolecular reaction, see: Balamurugan R, Gudla V. Synthesis of arylnaphthalene lignan scaffold by Gold-catalyzed intramolecular sequential electrophilic addition and benzannulation. *J Org Chem*, 2011, 76: 9919–9933

6 For the review, see: Yamamoto H, *Lewis Acids in Organic Synthesis*, John Wiley & Sons: New York, 2000, 96–113; For some other examples, see: (a) Pastine SJ, McQuaid KM, Sames D. Room temperature hydroalkylation of electron-deficient olefins: sp^3 C–H Functionalization via a Lewis acid-catalyzed intramolecular redox event. *J Am Chem Soc*, 2005, 127: 12180–12181; (b) McQuaid KM, Sames D. C–H bond functionalization via hydride transfer: Lewis acid catalyzed alkylation reactions by direct intramolecular coupling of sp^3 C–H bonds and reactive alkenyl oxocarbenium intermediates. *J Am Chem Soc*, 2009, 131: 402–403; (c) Pastine SJ, Sames D. Room temperature intramolecular hydro-O-alkylation of aldehydes: sp^3 C–H functionalization via a Lewis acid catalyzed tandem 1,5-hydride transfer/cyclization. *Org Lett*, 2005, 7: 5429–5431

7 Xiang SK, Wu G, Zhang B, Cui Y, Jiao N. Csp^2–Csp^2 bond formation via Lewis acid/ammonium salt cocatalyzed tandem addition and oxidative dehydrogenation strategy: alkenylation of indoles with α,β-unsaturated ketones. *Tetrahedron Lett*, 2012, 53(29): 3802–3804

8 Jia SX, Wang ML. Synthetic method of 3,4-dimethoxybenzene acetaldehyde. *Chem Indus Times*, 2006, 20(2): 21–22 (in Chinese)

SCIENCE CHINA
Chemistry

• SUPPORTING INFORMATION •


July 2013 Vol.56 No.7: 1–9
doi: 10.1007/s11426-013-4843-7

Synthesis of naphthalene derivatives through inexpensive BF_3·Et_2O-catalyzed annulation reaction of arylacetaldehydes with arylalkynes

XIANG ShiKai[1*], HU Hao[1], MA Jing[1], LI YuanZhuo[1], WANG BiQin[1*], FENG Chun[1], ZHAO KeQing[1], HU Ping[1] & CHEN XiaoZhen[2]


[1]*College of Chemistry and Materials Science, Sichuan Normal University, Chengdu 610066, China*
[2]*Chengdu Institute of Biology, Chinese Academy of Sciences, Chengdu 610041, China*


Received November 11, 2012; accepted December 26, 2012; published online February 21, 2013





*Corresponding authors (email: xiangsk@hotmail.com; wangbiqin1964@126.com)


© Science China Press and Springer-Verlag Berlin Heidelberg 2013



3aa
^{13}C NMR
140.716
140.215
133.744
131.571
130.044
128.230
127.595
127.199
126.900
125.992
125.729
125.352
77.318
77.000
76.682
200
150
100
50
0 PPM



Me
3ba
^{1}H NMR
7.803
7.783
7.657
7.480
7.458
7.432
7.417
7.375
7.361
7.322
7.301
2.419
3.99
1.90
0.91
1.70
3.00
10
9
8
7
6
5
4
3
2
PPM


Me
3ba
^{13}C NMR
140.958
139.529
135.704
132.000
131.706
130.037
128.215
128.100
127.983
127.357
127.090
127.037
124.819
124.452
77.319
77.000
76.682
21.932
200
150
100
50
PPM


MeO
3ca
^{1}H NMR
7.807
7.785
7.518
7.501
7.482
7.465
7.426
7.411
7.382
7.370
7.225
7.171
7.149
3.920
3.745
3.96
3.00
1.92
1.90
0.26
10
8
6
4
2
PPM



OMe
3da
^{1}H NMR
8.321
8.300
7.520
7.503
7.471
7.469
7.412
7.327
7.307
7.287
6.817
6.799
3.994
7.88
3.00
0.93
1.04
1.02
10
8
6
4
2
0PPM



OMe
3da
^{13}C NMR
155.538
141.093
139.809
130.056
132.598
128.136
127.512
127.095
125.897
125.816
124.660
121.476
118.324
103.627
77.318
77.000
76.682
55.570
200
150
100
50
PPM




3ea

^{1}H NMR











3ea

^{13}C NMR













3fa

^{1}H NMR













^{1}H NMR





^{13}C NMR



Bu^t 3ac
^{1}H NMR
7.971
7.949
7.906
7.885
7.850
7.830
7.530
7.513
7.498
7.482
7.462
7.446
7.441
7.436
7.432
7.420
7.402
1.409
9.00
4.11
3.56
0.93
0.93
0.92
10
8
6
4
2
0
PPM



Bu^t 3ac
^{13}C NMR
150.061
140.202
137.734
133.801
131.696
129.722
128.220
127.395
126.913
126.177
125.876
125.683
125.398
125.154
77.330
77.000
76.682
34.597
31.481
200
150
100
50
PPM



Br 3ad
^{1}H NMR
7.896
7.875
7.861
7.838
7.816
7.601
7.595
7.585
7.488
7.487
7.480
7.435
7.430
7.418
7.349
7.332
3.10
2.07
1.92
1.00
1.91
0.89
10
8
6
4
2
0 PPM



Br
3ad
13C NMR
139.595
138.866
133.741
131.667
131.403
131.298
128.340
127.982
126.836
126.238
125.897
125.608
125.328
121.441
77.317
77.000
76.683
200
150
100
50
PPM



F
3ae
1H NMR
7.902
7.882
7.841
7.516
7.499
7.482
7.465
7.431
7.425
7.420
7.384
7.368
7.183
7.162
7.145
3.12
2.04
2.02
2.01
1.00
1.07
10
8
6
4
2
PPM



F
3ae
13C NMR
163.456
161.012
136.629
139.108
136.586
133.746
131.572
131.497
128.300
127.776
126.980
126.135
125.816
125.724
125.319
115.262
115.049
77.317
77.000
76.681
200
150
100
50
PPM



7.869
7.864
7.832
7.810
7.756
7.736
7.544
7.527
7.506
7.483
7.464
7.453
7.441
7.437
7.423
7.411
7.393
7.354
7.349
7.330
7.325
0.000
Cl
3ga-1
^{1}H NMR
7.33
1.06
1.05
1.03
10
8
6
4
2
PPM


Cl
3ga-1
^{1}H NMR
7.869
7.864
7.832
7.810
7.756
7.736
7.544
7.527
7.506
7.483
7.464
7.453
7.441
7.437
7.423
7.411
7.393
7.354
7.349
7.330
7.325
7.33
1.00
1.06
1.05
1.03
7.9
7.8
7.7
7.6
7.5
7.4
7.3
PPM


Cl
3ga-1
^{13}C NMR
140.388
140.194
134.463
131.614
129.933
129.896
128.346
127.845
127.464
127.106
126.809
126.733
126.535
77.318
77.000
76.675
200
150
100
50
PPM


Cl
3ga-2
^{13}C NMR
143.472
139.506
135.968
131.408
131.033
129.610
129.362
128.656
128.595
128.160
127.236
126.698
125.629
125.366
77.317
77.000
76.683
200
150
100
50
PPM


7.852
7.832
7.801
7.778
7.764
7.752
7.744
7.661
7.480
7.472
7.448
7.425
7.405
7.396
7.373
7.362
7.355
7.343
7.333
7.306
7.288
7.259
7.237
7.226
7.214
2.501
2.005
0.000
Me
(a)
and
Me
(b)
3ha-1
3ha-2
^{1}H NMR
15.21
3.53
0.96
(a)
(b)
3.00
2.62
10
8
6
4
2
PPM


(c)
Me
and
Me
3ha-1
3ha-2
7.852
7.832
7.801
7.778
7.764
7.752
7.744
7.661
7.480
7.472
7.448
7.425
7.405
7.396
7.373
7.362
7.355
7.343
7.333
7.306
7.288
7.259
7.237
7.226
7.214
15.21
(c)
3.53
0.96
8.2
8.0
7.8
7.6
7.4
7.2
7.0
6.8
PPM